arXiv:1211.2431v1 [math.SG] 11 Nov 2012

NONGENERIC J-HOLOMORPHIC CURVES IN SYMPLECTIC 4-MANIFOLDS

DUSA MCDUFF

Abstract. This note discusses the structure of J-holomorphic curves in symplectic 4-manifolds (M,ω) when $J \in \mathcal{J}(\mathcal{S})$, the set of ω-tame J for which a fixed chain $\mathcal{S}$ of transversally intersecting embedded spheres of self-intersection ≤ -2 is J-holomorphic. Extending work by Biran (in *Invent. Math.* 1999), it shows that when (M,ω) is the blow up of a rational or ruled symplectic 4-manifold any homology class $A \in H_2(M;\mathbb{Z})$, with nonzero Gromov invariant and nonnegative intersection both with the spheres in $\mathcal{S}$ and with the exceptional classes other than A, has an embedded J-holomorphic representative for some $J \in \mathcal{J}(\mathcal{S})$. This result is a key step in some of the arguments in McDuff (*Journ. Topology* (2009)) on embedding ellipsoids, and also has applications to symplectic 4-orbifolds.

Contents

1. Introduction 2
1.1. Background 2
1.2. Main results 3
2. Results using Taubes–Seiberg–Witten theory 7
2.1. Review of J-holomorphic curve theory 7
2.2. The case $J \in \mathcal{J}(\mathcal{S})$ 10
2.3. Representing exceptional spheres 13
3. Geometric general position arguments 14
3.1. Patching procedures 14
3.2. General position arguments 19
4. Proofs of the main results 24
4.1. Proof of Theorem 1.2.1 24
4.2. Numerical arguments 33
4.3. Symplectic inflation 36
References 40

Date: November 7, 2012.
Key words and phrases. J-holomorphic curve, rational symplectic 4-manifold.
partially supported by NSF grant DMS 0905191.

1. Introduction

1.1. **Background.** Let (M,ω) be a closed symplectic 4-manifold. Write $\mathcal{E} \subset H_2(M;\mathbb{Z})$ for the set of classes that can be represented by exceptional spheres, i.e. symplectically embedded spheres with self-intersection -1. It follows from work of Taubes [Tau96] as extended by Li–Liu [LL95, LL01] that if Taubes' version of the Gromov invariant $\mathrm{Gr}(A)$ is nonzero and if $A \cdot E \geq 0$ for all $E \in \mathcal{E} \smallsetminus \{A\}$, then for generic ω-tame almost complex structure J on (M,ω) the class A is represented by an embedded J-holomorphic curve. We are interested in the representatives of A with respect to special J for which some nongeneric chains of spheres $\mathcal{S}$ are holomorphic. In McDuff [Mc09i, Mc09ii], we asserted that even in this case any class A that is disjoint from $\mathcal{S}$ would be represented by a smoothly embedded J-holomorphic connected curve for suitable J, because we can assume that J is generic away from $\mathcal{S}$. This result, though it does turn out to be correct, requires proof, that we supply in this paper. Similar results were obtained in Biran [B99] in the case when all the curves in $\mathcal{S}$ are spheres of self-intersection -2.

Let us first describe the kind of singular set $\mathcal{S}$ we consider.

Definition 1.1.1. *The set* $\mathcal{S} := C^S_1 \cup \cdots \cup C^S_s$ *in* (M,ω) *is called a* **singular set** *if it is at union of symplectically embedded curves of genus* $g(S_i)$ *in classes* $S_1, \dots, S_s$ *respectively such that*

- (i) $S_i^2 = -k_i \leq -2$;
- (ii) *the curves* $(C^S_i)_{1\leq i\leq s}$ *intersect transversally, in such a way that at each intersection point their tangent spaces are symplectically orthogonal.*

A symplectic form is said to be **adapted** *to* $\mathcal{S}$ *if* $\mathcal{S}$ *satisfies these conditions with respect to it. Each* $\mathcal{S}$ *defines a* **graph** $\Gamma_{\mathcal{S}}$ *with vertices* $v_1, \dots, v_s$*, and in which each pair* v_i, v_j *of distinct vertices are joined by precisely* $S_i \cdot S_j$ *edges.*

Given $\mathcal{S}$, we can identify a neighborhood $\mathcal{N}(C^S_i)$ of each C^S_i with a neighborhood of the zero section in a holomorphic line bundle $\mathcal{L}_{-k_i}$ over C^S_i $S^2 = \mathbb{C} \cup \{\infty\}$ with Chern class $-k_i$, and obtain $\mathcal{S}$ by plumbing, i.e. for each point $q \in C^S_i \cap C^S_j$ where $i \neq j$ we identify a neighborhood of $q \in C^S_i$ in $\mathcal{N}(C^S_i)$ with a neighborhood of $q \in C^S_j$ in $\mathcal{N}(C^S_j)$ preserving the local product structure but interchanging fiber and base. We call the resulting plumbed structure on the neighborhood $\mathcal{N}(\mathcal{S}) = \cup_i \mathcal{N}(C^S_i)$ the **local fibered structure**.

Definition 1.1.2. *We say that the singular set* $\mathcal{S}$ *is* **negative** *if for every tuple* $(m_1, \dots, m_s)$ *with* $m_i \geq 0$ *for all* i*, we have*

$$\Big(\sum_{i=1}^{s} m_i S_i\Big) \cdot S_j \geq 0 \ \forall j \Longrightarrow m_i = 0 \ \forall i.$$

Further, we say that $\mathcal{S}$ *(or* $\Gamma_{\mathcal{S}}$*) is a* **chain** *if the following conditions hold:*

- $C^S_i \cdot C^S_j \leq 1$ *for all* $i \neq j$*; and*
- C^S_i *is disjoint from* C^S_j *when* $|i-j| > 1$.

Note that we allow chains to be disconnected. If $\Gamma_{\mathcal S}$ is not a chain, then we will usually assume that it is at least a **forest**, i.e. a disjoint union of trees. Further we always assume that $k_i := -S_i \cdot S_i \ge 2$ for all i.

Lemma 1.1.3. *Every chain is negative.*

Proof. Consider a nonnegative and nonzero tuple $(m_1, \dots, m_s)$. If we ignore all i such that $m_i = 0$ then the singular set is still a chain. Further for each i there are at most two $j \ne i$ such that $S_i \cdot S_j \ne 0$, and there is i (with $m_i \ne 0$) with just one such j. Hence we have

$$\Big(\sum_i m_i S_i\Big) \cdot \Big(\sum m_j S_j\Big) = \sum_i m_i S_i \cdot S_i + \sum_i m_i S_i \Big(\sum_{j \ne i} S_j\Big) < -\sum_i m_i k_i + 2 \sum_i m_i \le 0.$$

Therefore this tuple does not satisfy the condition $\big(\sum_{i=1}^s m_i S_i\big) \cdot S_j \ge 0$ for all j. The result follows. □

Definition 1.1.4. *Given any singular set $\mathcal S$, we define $\mathcal J(\mathcal S) := \mathcal J(\mathcal S, \omega)$ to be the space of all ω-tame almost complex structures J on (M, ω) that are integrable in some neighborhood $\mathcal N(\mathcal S)$ and are compatible with its local fibered structure. Thus each C_i^S is J-holomorphic, as is each local projection $\mathcal N(C_i^S) \to C_i^S$. Moreover, if C_j^S intersects C_i^S the intersection $C_j^S \cap \mathcal N(C_i^S)$ coincides with a union of fibers of the projection $\mathcal N(C_i^S) \to C_i^S$.*

Example 1.1.5. (i) Suppose that (M, ω, J) is a toric manifold whose moment polytope has a connected chain of edges $\epsilon_i, i = 1, \dots, s$, with Chern numbers $-k_i$. Then the inverse image $\mathcal S$ of this chain of edges under the moment map is a chain of spheres with respect to the natural complex structure on M. Moreover the toric symplectic form is adapted to $\mathcal S$: in particular the S_i do intersect orthogonally. Another example of $\mathcal S$ is a disjoint union of embedded spheres each with self-intersection ≤ -2.

(ii) If $\mathcal S$ is a tree, then the condition $k_i \ge 2$ for all i is not enough to guarantee that $\mathcal S$ is negative. For example we could have $k_i = 2$ for $i = 1, \dots, 5$ and $S_i \cdot S_j = 1$ for $i = 1, j > 1$ and equal to 0 for all other pairs $i < j$. Then $B = 2S_1 + \sum_{i>1} S_i$ has $B \cdot S_i = 0$ for all i. The negativity of $\mathcal S$ is important in Proposition 1.2.5, but is not explicitly used in the proof of Theorem 1.2.1, though it is satisfied there because $\mathcal S$ is a chain.

1.2. Main results.

Recall that $\mathcal E$ denotes the set of classes that are represented by exceptional spheres. We often use the fact that $E \cdot E' \ge 0$ for all $E, E' \in \mathcal E$. This holds because for sufficiently generic J each $E \in \mathcal E$ has an embedded J-holomorphic representative.

Theorem 1.2.1. *Let (M, ω) be a blow up of a rational or ruled manifold, and suppose that the singular set $\mathcal S = C_1^S \cup \dots \cup C_s^S$ is a chain. Let $A \in H_2(M)$ be a nonzero class such that*

(i) $\mathrm{Gr}(A) \ne 0$*;*
(ii) *if $A^2 = 0$ then A is a primitive class;*

(iii) $A \cdot E \geq 0$ *for every* $E \in \mathcal{E} \smallsetminus \{A\}$*; and*
(iv) $A \cdot S_i \geq 0$ *for* $1 \leq i \leq s$.

Then there is a smooth embedded connected curve C^A *in class* A *that is* J*-holomorphic for some* $J \in \mathcal{J}(\mathcal{S})$ *and intersects each* C_i^S *positively and transversally in precisely* $S_i \cdot A$ *points.*

Note that we did not include classes $A = mA_0$ with $A^2 = 0$ and $m > 1$ in the statement above, since we wanted to assert that A has a connected representative. But if we apply the Theorem 1.2.1 to the underlying primitive class A_0 then we find that such A will be represented by the union of m disjoint copies of an embedded A_0-curve. Here is a translation of condition (i) in the case when M is rational, i.e. $S^2 \times S^2$ or a blowup of $\mathbb{CP}^2$.

Corollary 1.2.2. *If* (M, ω) *is rational, then the conclusion of Theorem 1.2.1 holds for any class* A *that satisfies (iii), (iv), has* $\omega(A) > 0$ *and* $d(A) := A^2 + c_1(A) \geq 0$*, and is such that one of the following conditions holds:*
(a) $A^2 > 0$*;*
(b) $A^2 = 0$ *and* A *is a primitive class;*
(c) $A \in \mathcal{E}$.

The proof that Corollary 1.2.2 follows from Theorem 1.2.1 is given in Lemma 2.1.3. There we also discuss the analogous conditions for blow ups of ruled surfaces.

Corollary 1.2.3. *If* $\Gamma_{\mathcal{S}}$ *is a chain and a class* A *as in Theorem 1.2.1 can be represented by a symplectically embedded curve of genus* $g(A)$*, then the components of any stable map representing* A *that is* J*-holomorphic for a generic element in* $\mathcal{J}(\mathcal{S})$ *are parametrized by curves whose genera sum to at most* $g(A)$*. Moreover, this inequality is strict if any component is not embedded or if there is a pair of components that intersect more than once.*

Remark 1.2.4. (i) Note that Corollary 1.2.3 is not immediately obvious since we cannot assume that an arbitrary J-holomorphic stable map in class A is the limit of embedded curves, even for generic elements $J \in \mathcal{J}(\mathcal{S})$. Our proof does use the fact that J is generic, though the recent preprint Li–Zhang [LZ12] shows by purely algebraic methods that the first claim holds for arbitrary $J \in \mathcal{J}(\mathcal{S})$ provided that A is J-NEF, in other words that it has nonnegative intersection with every class that has a J-holomorphic representative. This paper also explains other severe restrictions on the intersection patterns of these curves in the case when A is J-NEF and $g(A) = 0$. However, these results are not as helpful to us as one might think, since sometimes (as in the proof of Proposition 1.2.5) the essence of our argument is to show that A is J-NEF.

Intuitively, the statement in Corollary 1.2.3 is clear. For example if $A \cdot S_i = 0$ for all i, then an embedded representative of A can be considered as a smoothing of the singular curve in the (singular) space obtained by blowing down the components of $\mathcal{S}$; but, in general, smoothing a singularity creates genus. Thus if the smoothed curve has genus $g(A)$ the components of its nodal representative must also have genus $\leq g(A)$.

(ii) Our proof of Theorem 1.2.1 does not work if $\mathcal{S}$ has a component S_i with $S_i^2 = -1$, or if the graph $\Gamma_{\mathcal{S}}$ has a vertex of degree > 2: see Remark 4.1.5. However, as we explain there it seems likely that our methods can be significantly extended. This is the subject of ongoing research.

(iii) The restriction of Theorem 1.2.1 to the case of blowups of rational and ruled manifolds is mostly a matter of convenience; many of our arguments are geometric and hence are valid for any 4-manifold. However, some of the assertions (for example that the curve C^A is connected) use Lemma 2.1.1 and hence are valid only when $b_2^+ = 1$. Since our main purpose in proving Theorem 1.2.1 is to use it to deduce Proposition 1.2.9 about the method of inflation relative to $\mathcal{S}$, and since this works only in the case of rational and ruled manifolds and their blowups, we have restricted to that case.

(iv) Our constructions start from a J_0-holomorphic stable map (or nodal curve) Σ^A representing the class A, where $J_0 \in \mathcal{J}(\mathcal{S})$ is assumed "generic" outside $\mathcal{S}$. (More precisely, J lies in the set $\mathcal{J}_{reg}(\mathcal{S}, A, \omega)$ described in Definition 2.2.1.) The resulting embedded curve C^A is J-holomorphic for some $J \in \mathcal{J}_{reg}(\mathcal{S}, A, \omega)$ that equals J_0 near $\mathcal{S}$, and otherwise is $\mathcal{C}^1$-close to J_0 and equal to it outside a neighborhood of the image of the original stable map; cf. Remark 3.1.4.

When $A \in \mathcal{E}$, we can make a much stronger statement. It applies for all M, not just those with $b_2^+ = 1$.

Proposition 1.2.5. *Let (M, ω) be any symplectic 4-manifold, and $\mathcal{S}$ be any negative singular set, for example a chain. Suppose further that the class A in Theorem 1.2.1 lies in $\mathcal{E}$. Then there is a path connected and residual subset $\mathcal{J}_{SW}(\mathcal{S}, A)$ of $\mathcal{J}(\mathcal{S})$ such that for each $J \in \mathcal{J}_{SW}(\mathcal{S}, A)$ the only J-holomorphic representative of the class A is an embedded sphere C^A.*

This is proved at the end of §2.3. The proof does not need the geometric techniques developed for Theorem 1.2.1; rather it exploits the fact that $A^2 < 0$.

Remark 1.2.6. In the above proposition, we take $\mathcal{J}(\mathcal{S})$ as in Definition 1.1.4. However, the result still holds if we look at the subset of $\mathcal{J}(\mathcal{S})$ consisting of elements that restrict to a fixed element J_0 on some, arbitrarily small, neighborhood of $\mathcal{S}$. For example, if the components of $\mathcal{S}$ have a local toric structure as in Example 1.1.5 then we might take J_0 to be the corresponding complex structure.

For other classes A, although Theorem 1.2.1 asserts the existence of embedded representatives C^A for some $J \in \mathcal{J}(\mathcal{S})$, our geometric arguments do not show that these exist for an arbitrary generic[1] $J \in \mathcal{J}(\mathcal{S})$, even in the case of spheres. However, one can show this in certain cases by numerical (dimension counting) arguments. In §4.2 we first explain an argument due to Li–Zhang [LZ12] that applies to spherical classes, i.e. when $g(A) := 1 + \frac{1}{2}(A^2 - c_1(A)) = 0$. Secondly, we extend an argument due to Biran [B99], that applies when the curves in $\mathcal{S}$ are "not too singular." We then prove the following result in Corollary 4.2.4. Here $d(A) := c_1(A) + A^2$ is the Seiberg–Witten degree; cf. §2.1.[2]

---

[1] More precisely, to say that a given property holds for a generic $J \in \mathcal{J}(\mathcal{S})$ means that there is a residual subset $\mathcal{J}'(\mathcal{S})$ of $\mathcal{J}(\mathcal{S})$ on which it holds.

Proposition 1.2.7. *Let (M, ω) be any symplectic 4-manifold with $b_2^+ = 1$, and let $\mathcal{S}$ be a singular set whose components are spheres with $c_1(S_i) = 0$ but with any graph $\Gamma_{\mathcal{S}}$. Suppose further that A satisfies the conditions of Theorem 1.2.1. Then the following holds.*

- (i) *If, in addition, A cannot be written as $\sum_i \ell_i S_i$ for $\ell_i \geq 0$, then A is represented by an embedded J-holomorphic curve C^A for generic $J \in \mathcal{J}(\mathcal{S})$. In fact there is at least one such curve through each generic set of $\frac{1}{2}d(A)$ points in (M, ω).*
- (ii) *If $\Gamma_{\mathcal{S}}$ is a forest, then A is never equal to $\sum_i \ell_i S_i$ for $\ell_i \geq 0$.*

Moreover, the conclusion in (i) holds for any negative singular set $\mathcal{S}$ provided that A satisfies the additional condition that $g(A) = 0$.

Remark 1.2.8. (i) Biran [B99, Lemma 2.2B] proved the special case of Theorem 1.2.1 and Proposition 1.2.5 when $\mathcal{S}$ is a union of disjoint spheres of self-intersections $-k_i \leq -2$ and $A \cdot S_i + c_1(S_i) \geq 0$ for all i. In [Mc09i] we are mostly interested in the case when $A \cdot S_i = 0$ for all i. In this case, Biran's argument applies when $k_i = 2$ for all i. The proof of Lemma 4.2.1 shows that Biran's argument extends to the case when the spheres S_i have arbitrary (transverse) intersections, but still $k_i = 2$ for all i. This is enough to fill in all gaps in [Mc09ii]; cf. Remark 4.2.5. However, it does not suffice for [Mc09i] where the k_i can be arbitrary. In this case, one seems to need geometric arguments in order to prove Theorem 1.2.1.

(ii) Even though there is considerable overlap between Proposition 1.2.5 and the last claim in Proposition 1.2.7 about the genus 0 case, there are subtle differences both in the statements and the proofs. In particular, Proposition 1.2.5 applies to any 4-manifold, not just those with $b_2^+ = 1$, but restricts to the rigid case $A \in \mathcal{E}$. See also Remark 4.2.5 (iii).

As in Biran [B99], the main application of Theorem 1.2.1 is to show that certain symplectic inflation processes are possible: one uses the curve C_A to construct a form ρ_A that represents the Poincaré dual to A by means of which one can deform the initial symplectic form. For this, it is enough to have a single curve. However, [Mc09i, Mc09ii] also applies the method of symplectic inflation from McDuff [Mc98] to conclude that certain embedding spaces are connected; cf. [Mc09i, Corollary 1.6]. To prove this one needs embedded J-holomorphic representatives of A for 1-parameter families of J. We construct these in §4.3; cf. Lemma 4.3.3. The following relative version of "deformation implies isotopy" is a typical corollary.

Proposition 1.2.9. *Let (M, ω) be a blow up of a rational or ruled 4-manifold, and $\mathcal{S} \subset M$ be a chain of ω-orthogonally intersecting spheres as above. Let ω' be any symplectic form on M such that*

---

[2] For simplicity, we restrict to the case $b_2^+ = 1$. If $b_2^+ > 1$, there is an analogous statement. However, in that case the hypothesis $\mathrm{Gr}(A) \neq 0$ implies that $d(A) = 0$, so that the result is less interesting.

(a) $[\omega'] = [\omega] \in H^2(M)$;
(b) $\omega' = \omega$ *in some neighborhood of* $\mathcal{S}$;
(c) *there is a family of noncohomologous symplectic forms* $\omega_t, t \in [0,1]$, *on* M *that equal* ω *near* $\mathcal{S}$ *and are such that* $\omega_0 = \omega$ *and* $\omega_1 = \omega'$.

Then there is a compactly supported isotopy ϕ_t *of* $M \smallsetminus \mathcal{S}$ *such that* $\phi_0 = id$ *and* $\phi_1^*(\omega) = \omega'$.

Acknowledgements. I warmly thank E. Opshtein for noticing and discussing the gap in the proofs in [Mc09i], and Matthew Strom Borman, Tian-Jun Li and Felix Schlenk for very helpful comments on earlier drafts of this paper.

2. Results using Taubes–Seiberg–Witten theory

This section collects the results that can be proved by modifications of standard J-holomorphic curve theory in dimension 4. In particular, we prove Proposition 1.2.5 in §2.2. We begin by recalling relevant parts of this theory. Throughout, unless specific mention is made to the contrary,[3] by a **curve** we mean the image of a smooth map $u : \Sigma \to M$ where Σ is a connected smooth Riemann surface. Thus an **immersed J-holomorphic curve** is the image of a smooth J-holomorphic immersion $u : \Sigma \to M$. In particular, all its double points have positive intersection number. A curve is called **simple** (or **somewhere injective**) if it is not multiply covered; cf. [MS04, Chapter 2].

2.1. Review of J-holomorphic curve theory.

We begin this section by a very brief review of Taubes' work relating Seiberg–Witten theory to J-holomorphic curves in order to explain the first hypothesis in Theorem 1.2.1, namely that $\mathrm{Gr}(A) \neq 0$. Here, $\mathrm{Gr}(A)$ is Taubes' version of the Gromov invariant of A, that to a first approximation counts embedded J-holomorphic curves in (M,ω) through $\frac{1}{2}d(A)$ generic points, where $d(A) := c_1(A) + A^2$ is the index of the appropriate Fredholm problem; cf. [Tau96]. Thus $\mathrm{Gr}(A) \neq 0$ implies both that $d(A) \geq 0$ and that $\omega'(A) > 0$ for all symplectic forms ω' that can be joined to ω by a deformation (i.e. a path of possibly noncohomologous symplectic forms). For 4-manifolds with $b_2^+ = 1$ (such as blow ups of rational and ruled manifolds), one shows that $\mathrm{Gr}(A) \neq 0$ by using the wall crossing formulas in Kronheimer–Mrowka [KM94] in the rational case and Li–Liu [LL95] in the ruled case.

Note first that because the intersection form on $H^2(M;\mathbb{R})$ has type $(1,N)$, the cone $\mathcal{P} := \{a \in H^2(M) \mid a^2 > 0\}$ has two components; let $\mathcal{P}^+$ be the component containing $[\omega]$. Then we have the following useful fact.

Lemma 2.1.1. *Suppose that* $b_2^+(M) = 1$. *If* $a, b \in \overline{\mathcal{P}^+} \smallsetminus \{0\}$ *then* $a \cdot b \geq 0$ *with equality only if* $a^2 = 0$ *and* b *is a multiple of* a.

Taubes' Gromov invariant $\mathrm{Gr}(A)$ in [Tau96] counts holomorphic submanifolds and hence is somewhat different from the "usual" invariant due to Ruan–Tian that counts (perturbed) J-holomorphic maps $u : (\Sigma, j) \to (M, J)$ with a connected domain of fixed topological type modulo reparametrization. To explain the relation, we first make the following definition.

---

[3] Occasionally we allow a curve to be disconnected, but it never has nodes unless the adjective "nodal" or "stable" is used.

Definition 2.1.2. *We say that a class $A \in H_2(M;\mathbb{Z})$ is* **reduced** *if $A \cdot E \geq 0$ for all $E \in \mathcal{E} \smallsetminus \{A\}$.*

For example, as noted at the beginning of §1.2, every $E \in \mathcal{E}$ is reducd. If A is reduced and has $d(A) \geq 0$, then the adjunction formula $c_1(A) = 2-2g+A^2$ for embedded curves determines a unique genus $g(A)$ for which A can be represented by a holomorphic submanifold. Moreover, positivity of intersections ([Mc94i] or [MS04, Appendix E]) implies that any J-holomorphic map $(\Sigma_{g(A)}, j) \to (M, J)$ with $g = g(A)$ must be an embedding, since any singular point gives rise to a defect in the genus formula. It follows that, except in the case of tori of zero self-intersection (where double covers affect the count in a very delicate way), the invariant $\mathrm{Gr}(A)$ for reduced A simply counts (with appropriate signs) the number of possibly disconnected J-holomorphic curves of genus $g(A)$ through $\frac{1}{2}d(A)$ generic points. For example, $\mathrm{Gr}(E) = 1$ for all $E \in \mathcal{E}$. In fact, one can check that the only reduced classes A with $A^2 < 0$ and $d(A) \geq 0$ are those of the exceptional spheres $A \in \mathcal{E}$.

If, for a reduced A, the Gromov invariant $\mathrm{Gr}(A)$ counts disconnected curves, then A must decompose as a sum $A_1 + A_2$ with $A_1 \cdot A_2 = 0$ and $\mathrm{Gr}(A_1), \mathrm{Gr}(A_2)$ both nonzero. Further, $A_i^2 \geq 0$ for $i = 1, 2$, since otherwise, as we saw above $A_i \in \mathcal{E}$ and then $A \cdot A_i = A_i^2 < 0$ implies that A is not reduced. But then, when $b_2^+(M) = 1$, we have $A_i \in \overline{\mathcal{P}}^+$ for $i = 1, 2$ which implies by Lemma 2.1.1 that A, A_1, A_2 all have square 0 and lie on the same line. Hence every reduced class with $A^2 > 0$ is represented by a connected curve. This is also true when $A^2 = 0$ unless A is not primitive, that is $A = mA_0$ for some $m > 1$. In that case, the adjunction formula implies that either $g(A_0) = 0$, $d(A) = 2m$ and A is represented by parallel copies of A_0 through m generic points, or that $g(A_0) = 1$ and $d(A) = 0$, in which case the counting procedure when $m > 1$ takes double covers into account and so is more complicated (see [Tau96]). In sum, we have:

> *When $b_2^+(M) = 1$ for reduced classes A that are not represented by embedded tori of zero self-intersection,* $\mathrm{Gr}(A)$ *equals the standard J-holomorphic curve invariant for curves with genus $g(A)$.*

The situation when $b_2^+(M) > 1$ is more complicated, but we will not need this case.

Now let us consider a general, not necessarily reduced class, A with $\mathrm{Gr}(A) \neq 0$. Then it is shown in McDuff [Mc97, Proposition 3.1] that if we decompose A as

$$A = B + \sum_{E \in \mathcal{E}(A)} |A \cdot E|\, E, \qquad \mathcal{E}(A) = \{E \mid E \cdot A < 0\},$$

then $E \cdot E' = 0$ for $E, E' \in \mathcal{E}(A)$ and B is reduced with $d(B) \geq d(A) \geq 0$ and $B \cdot E = 0$ for all $E \in \mathcal{E}(A)$. Thus, for generic J the class A is represented by a main (possibly empty) embedded component C^B in class B, together with a finite number of disjoint curves C^E each with multiplicity $|A \cdot E|$ in the classes $E \in \mathcal{E}(A)$. This is proved by considering the structure of a J-holomorphic A-curve (where J is generic) through $\frac{1}{2}d'(A)$ points, where

$$d'(A) = c_1(A) + A^2 + \sum_{E\in\mathcal{E}} \big(|A\cdot E|^2 - |A\cdot E|\big). \tag{2.1.1}$$

It follows that $d'(A) = d(B)$. Moreover, Li–Liu show in [LL99] that the equivalence between Seiberg–Witten and Gromov invariants, previously established for reduced classes, extends to show that the class A has the same invariants as does its reduction B. Thus (except when there are multiply covered tori) $\mathrm{Gr}(A) = \mathrm{Gr}(B)$ counts the number of embedded B-curves through $d'(A) = d(B)$ generic points. In particular, if $B = 0$ then $d'(A) = d(B) = 0$.

We next discuss conditions that imply $\mathrm{Gr}(A) \neq 0$. The following is a sharper version of Li–Liu [LL01, Proposition 4.3]. (Their result applies to more general manifolds.)

Lemma 2.1.3. (i) *Let (M,ω) be a blowup of $\mathbb{C}P^2$. If $A \in H_2(M)$ satisfies $A^2 \geq 0$, $\omega(A) > 0$, and $d(A) \geq 0$, then* $\mathrm{Gr}(A) \neq 0$. *Further, Corollary 1.2.2 holds.*

(ii) *Let M be the k-point blowup of a ruled surface with base of genus $g(M) \geq 1$. Then a sufficient (but not necessary) condition for* $\mathrm{Gr}(A)$ *to be nonzero is that $A \in \mathcal{P}^+$ and $d(A) > g(M) + \frac{k}{4}$.*

(iii) *Let M be as in (ii) and $A \in H_2(M)$ be in the image of the Hurewicz map $\pi_2(M) \to H_2(M)$. Then $d(A) \geq 0$ implies that* $\mathrm{Gr}(A) \neq 0$.

Proof. We prove (i). By hypothesis (M,ω) is obtained from the standard $\mathbb{C}P^2$ by blowing up $N \geq 0$ points. Let $E_i \in H_2(M), i = 1,\dots,N$, be the classes of the corresponding exceptional divisors. Then the anticanonical class $K = -c_1(M)$ is standard, namely $K = -3L + \sum_{i=1}^N E_i$, where $L = [\mathbb{C}P^1]$. (As usual we identify $H_2(M;\mathbb{Z})$ with $H^2(M;\mathbb{Z})$ via Poincaré duality.) Because $d(A) \geq 0$, it follows from the wall crossing formula in [KM94] that exactly one of $\mathrm{Gr}(A), \mathrm{Gr}(K-A)$ is nonzero. Since $A^2 \geq 0$ and $\omega(A) > 0$, the Poincaré dual of A lies in $\overline{\mathcal{P}^+}$. Hence Lemma 2.1.1 implies that $\omega'(A) > 0$ for all forms ω' obtained from ω by deformation. On the other hand if $\omega'(E_i)$ is sufficiently small for all i, $\omega'(K-A) < 0$. Therefore $K - A$ has no J-holomorphic representative for ω'-tame J, so that $\mathrm{Gr}(K-A)$ must be zero. Hence $\mathrm{Gr}(A) \neq 0$. This proves the first claim in (i).

To prove Corollary 1.2.2 we must check that if A satisfies (a), (b) or (c) then conditions (i) and (ii) in Theorem 1.2.1 hold. The argument above proves this when A satisfies (a) or (b). Case (c) concerns the case $A \in \mathcal{E}$. But by standard Gromov theory, these classes have unique J-holomorphic representatives for all ω-tame and generic J, which again implies that $\omega'(E) > 0$ for all ω' deformation equivalent to ω. Hence as above, we must have $\mathrm{Gr}(A) \neq 0$ in this case as well. This completes the proof of (i).

To prove (ii), we use [LL01, Lemma 3.4] which states that $\mathrm{Gr}(A) \neq 0$ if $d(A) \geq 0$ and $2A - K \in \mathcal{P}^+$. Therefore (ii) will hold provided that $2A - K \in \mathcal{P}^+$. Suppose that $M = (S^2 \times \Sigma_g)\#k\mathbb{C}P^2$ is the k-fold blow up of the trivial bundle, where the exceptional divisors are $E_1,\dots,E_k$. Then $K = -2[\Sigma_g] + 2(g-1)[S^2] + \sum_{i=1}^k E_i$ so that $K^2 = -8(g-1) - k \leq 0$. Hence the nontrivial ruled surface over Σ_g also has

$K^2 = -4(g-1)$, since its one point blowup is the same as the one point blowup of the trivial bundle and blowing up reduces K^2 by 1. Then

$$(2A-K)^2 = 4A^2 - 4A\cdot K + K^2 = 4d(A) - 4(g-1) - k > 0,$$

by our assumption. Therefore either $2A-K \in \mathcal{P}^+$ or $-(2A-K) \in \mathcal{P}^+$. But the displayed inequality also shows that $A\cdot K \le A^2 + \frac{1}{4}K^2 \le A^2$ so that $A\cdot(2A-K) = 2A^2 - A\cdot K > 0$. Hence, because $A \in \mathcal{P}^+$, Lemma 2.1.1 implies that $2A-K \in \mathcal{P}^+$ as required.

To prove (iii), let us first consider the case when (M,ω) is minimal. Then $A = k[S^2]$, where $[S^2]$ is the class of the fiber. Further $k \ge 1$ since $d(A) = c_1(A) \ge 0$. Hence $\mathrm{Gr}(A) := \mathrm{Gr}(M,A) \ne 0$ by direct calculation. Note that this class takes values in $\mathbb{Z} \equiv \Lambda^0 H^1(M;\mathbb{Z})$. We can now use the blow down formula of [LL01, Lemma 2.8]. This says that if (X,τ) is obtained from (X',τ') by blowing down the single exceptional class E, and if $\mathrm{Gr}(X,A)$ takes values in $\mathbb{Z} \equiv \Lambda^0 H^1(X;\mathbb{Z})$, then for all $\ell \ge 0$

$$d(B-\ell E) \ge 0 \Longrightarrow \mathrm{Gr}(X', B-\ell E) = \mathrm{Gr}(X,B) \in \mathbb{Z}.$$

Note finally that $d(B-\ell E) = (B-\ell E)^2 + c_1(B) + \ell = d(B) - \ell(\ell-1) \le d(B)$. Therefore if we start with a class in the k-fold blow up with $d(A) \ge 0$, as we blow it down the degree $d(A)$ increases and we end up with a class $k[S^2], k > 0$, in the underlying minimal ruled surface. Hence $\mathrm{Gr}(A) \ne 0$. □

Finally, the following recognition principle will be useful. It is taken from [MS96, Corollary 1.5], but here we explain some extra details in the proof.

Lemma 2.1.4. *Suppose that (M^4,ω) admits a symplectically embedded submanifold with $c_1(Z) > 0$ that is not an exceptional sphere. Then (M,ω) is the blow up of a rational or ruled manifold.*

Proof. Since $0 < c_1(Z) = 2 - 2g + Z^2$, where g is the genus of the submanifold Z, we must have $Z^2 \ge 0$, since otherwise $Z^2 = -1$ and $g = 0$ so that Z is an exceptional sphere. But when $Z^2 \ge 0$ we can use the method of symplectic inflation from [L94] (cf. §**??** below) to deform ω to a symplectic form in class $[\omega_\kappa] := [\omega] + \kappa \mathrm{PD}(Z)$ for any $\kappa \ge 0$. Therefore if K is Poincaré dual to $-c_1(M)$, then $K\cdot Z < 0$ so that for large κ we have $\omega_\kappa(K) < 0$. But by Taubes' structure theorems in [Tau95], this is impossible when $b_2^+ > 1$. Thus $b_2^+ = 1$. The rest of the proof now follows the arguments given in [MS96]. The crucial ingredient is Liu's result that a minimal manifold with $K^2 < 0$ is ruled. □

2.2. **The case $J \in \mathcal{J}(\mathcal{S})$.** When we consider J-holomorphic representatives for a reduced class A when $J \in \mathcal{J}(\mathcal{S},\omega)$, the situation is rather different since the curves in $\mathcal{S}$ are not regular. Thus A could decompose as $A = \sum_i \ell_i S_i + A'$ where $\ell_i \ge 0$, and we need to consider generic representations of the class A'. But A' need not be reduced, and hence could have a disconnected representative as above with some multiply covered exceptional spheres. As usual we write $\mathcal{J}(M,\omega)$ for the set of ω-tame almost complex structures, and $\mathcal{J}_{reg}(M,\omega)$ to be the subset of regular elements, i.e.

elements such that the linearized Cauchy–Riemann operator $D_{u,J}$ is surjective for all J-holomorphic maps $u : (\Sigma, j_\Sigma) \to (M, J)$. Here (Σ, j_Σ) is a smooth connected Riemann surface, and when $genus(\Sigma) > 0$ we allow the complex structure j_Σ on Σ to vary, so that the tangent space $T_{j_\Sigma}\mathcal{T}$ at j_Σ to Teichmüller space $\mathcal{T}$ is part of the domain of $D_{u,J}$; cf. [Mc97, Tau96]. Standard theory (as in [MS04, Chapter 3]) implies that $\mathcal{J}_{reg}(M, \omega)$ is residual in $\mathcal{J}(M, \omega)$.

Definition 2.2.1. *Suppose that A is reduced, so that $g(A)$ is defined. We then define the space $\mathcal{J}_{reg}(\mathcal{S}, \omega, A) =: \mathcal{J}_{reg}(\mathcal{S})$ of almost complex structures that are* **A-regular with respect to** *$\mathcal{S}$ to be the set of all $J \in \mathcal{J}(\mathcal{S}, \omega, A)$ such that the linearized Cauchy–Riemann operator $D_{u,J}$ described above is surjective for all somewhere injective elements u of the moduli space of J-holomorphic maps $\mathcal{M}(B, J) := \{u : (\Sigma, j_\Sigma) \to (M, J)\}$ in class B, where B satisfies the following conditions:*

(i) *$0 < \omega(B) \le \omega(A)$,*
(ii) *B is reduced, i.e. $B \cdot E \ge 0$ for all $E \in \mathcal{E} \setminus \{B\}$,*
(iii) *$B \cdot S_i \ge 0$ for all $S_i \in \mathcal{S}$.*

We further define the space $\mathcal{J}_{SW}(\mathcal{S}, \omega, A) =: \mathcal{J}_{SW}(\mathcal{S})$ to consist of all $J \in \mathcal{J}(\mathcal{S}, \omega, A)$ such that $c_1(B) > 0$ for all classes B that satisfy (i) and (iii) above and are represented by a J-holomorphic map $u : (S^2, j) \to (M, J)$.

Remark 2.2.2. Notice that if $u : (S^2, j) \to (M, J)$ is a somewhere injective J-holomorphic map in class B with $c_1(B) > 0$ then the Seiberg-Witten index $d(B)$ is also nonnegative. For $d(B) = c_1(B) + B^2$ and the adjunction formula $g(S^2) = 0 \le 1 + \frac{1}{2}(B^2 - c_1(B))$ implies that $B^2 \ge c_1(B) - 2 \ge -1$ with equality exactly if u is an embedding. Further, if $d(B) = 0$ then the only possibility is that $B^2 = -1$ and u is an embedding.

Lemma 2.2.3. *Suppose that $\Gamma_{\mathcal{S}}$ is negative. Then the following hold.*

(i) *The subset $\mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ of $\mathcal{J}(\mathcal{S})$ is residual in the sense of Baire.*
(ii) *$\mathcal{J}_{reg}(\mathcal{S}, \omega, A) \subset \mathcal{J}_{SW}(\mathcal{S}, \omega, A)$.*
(iii) *$\mathcal{J}_{SW}(\mathcal{S}, \omega, A)$ is path connected. Further, if $\omega_t, t \in [0, 1]$ is any smooth path of adapted symplectic forms, then the set*

$$\bigcup_{t \in [0,1]} \mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)$$

is path connected.

Proof. Since each $J \in \mathcal{J}(\mathcal{S})$ is fixed on some fibered neighborhood $\mathcal{N}(\mathcal{S})$ of $\mathcal{S}$ we first check that there is no class B satisfying the conditions (i), (ii) and (iii) in Definition 2.2.1 that has a J-holomorphic representative lying entirely in $\mathcal{N}(\mathcal{S})$. To see this, fix some such neighborhood $\mathcal{N}_0(\mathcal{S})$ and choose a map $\Phi : \mathcal{N}_0(\mathcal{S}) \to \mathcal{S}$ that agrees with the given fibration over some nonempty open subset of each curve C_i^S in $\mathcal{S}$. Then, for any J-holomorphic B-curve $u : \Sigma \to \mathcal{N}_0(\mathcal{S})$ the composite map $\Phi \circ u : \Sigma \to \mathcal{S}$ is J-holomorphic over an open subset of each C_i^S and hence must have nonnegative degree on each component. Thus $B := \sum_i m_i S_i$ where $m_i \geq 0$ for each i. Since $B \cdot S_j \geq 0$ by hypothesis on B and $\mathcal{S}$ is negative, we must have $B = 0$, contradicting (i).

It follows that each J-holomorphic B-curve must intersect $M \smallsetminus \mathcal{N}_0(\mathcal{S})$. By standard theory as in [MS04, Chapter 3.2], in order to ensure that every J-holomorphic B-curve is regular, it suffices to perturb J on $M \smallsetminus \mathcal{N}_0(\mathcal{S})$. One can then see that $\mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ is residual when $g(A)$ (and hence $g(B)$) is zero by arguing as in the proof of Theorem 3.1.5 [MS04]. One main technical ingredient is the version of the Riemann–Roch theorem in [MS04, Theorem C.1.10]. Since this is carried out for arbitrary genus, one can easily adapt the above proof to higher genus curves as in [Tau96, Mc97]. Since $\mathcal{J}_{reg}(\mathcal{S}, \omega, A) \subset \mathcal{J}_{SW}(\mathcal{S}, \omega, A)$, this proves (i).

To prove (ii), first recall that the (adjusted) Fredholm index $\mathrm{Ind}(D_{u,J})$ of a somewhere injective J-holomorphic curve $u : (S^2, j) \to (M^{2n}, J)$ in class B is $2n + 2c_1(B) - 6 = 2(c_1(B) - 1)$ in dimension $2n = 4$. (This is the virtual dimension of the quotient space of J-holomorphic maps modulo the action of the reparametrization group $G = PSL(2, \mathbb{C})$.) Since $\mathrm{Ind}(D_{u,J}) \geq 0$ for regular curves, such curves must lie in classes B with $c_1(B) > 0$. Thus (ii) holds.

Part (iii) follows by a similar argument. Consider a generic path $J_t \in \mathcal{J}(\mathcal{S}), t \in [0, 1]$, with endpoints in $J_0, J_1 \in \mathcal{J}_{SW}(\mathcal{S}, \omega, A)$, and let $u : S^2 \to M$ be J_t-holomorphic for some t. By standard Fredholm theory, $\dim \mathrm{Coker}\,(D_{u,J_t}) \leq 1$. Therefore, because $\mathrm{Ind}(D_{u,J})$ is even, every J_t-holomorphic map $u : S^2 \to M$ must have $\mathrm{Ind}(D_{u,J}) \geq 0$. Hence $c_1(B) \geq 0$ as in (ii). □

Lemma 2.2.4. (i) *If $d(B) \geq 0$ and $u : (\Sigma, j) \to M$ is an embedded J-holomorphic curve in class B for some $J \in \mathcal{J}(\mathcal{S}, \omega, A)$, then there is $J' \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ such that u is also J'-holomorphic.*

(ii) *If B with $\omega(B) \leq \omega(A)$ is represented by a connected J-homomorphic curve for some $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ and $B \neq S_i$ for any i, then $d(B) \geq 0$. Moreover, $B^2 \geq 0$ unless $B \in \mathcal{E}$, and if $B^2 = 0$ then B is represented by an embedded J-holomorphic sphere or torus.*

Proof. When $C^B := \mathrm{im}\, u$ is embedded, the pullback tangent bundle $(u^*(TM), J)$ splits as the sum of the tangent bundle $T\Sigma_g$ to the domain and the normal bundle. Further, the index of the linearization $D_{u,J}$ restricted to the normal line bundle to C_B is $d(B)$ and so nonnegative. Hence, by Riemann–Roch [MS04, Theorem C.1.10], we can adjust J in these normal directions to make $D_{u,J}$ surjective. Then the curve u itself is regular. Since every other J-holomorphic curve goes through points in $M \smallsetminus (\mathcal{S} \cup C^B)$, standard regularity arguments imply that by changing J outside a neighborhood of $\mathcal{S} \cup C^B$ we can arrange that J belongs to $\mathcal{J}_{reg}(\mathcal{S}, \omega, A)$. This proves (i).

To prove (ii), we use the fact that the the (adjusted) Fredholm index $\mathrm{Ind}(D_{u,J})$ of a somewhere injective J-holomorphic curve $u : (\Sigma_g, j) \to (M^4, J)$ in class B and with domain of genus g is $4(1 - g) + 2c_1(B) + 6g - 6 = 2(c_1(B) + g - 1)$, where as in the proof of Lemma 2.2.3 we adjust by quotienting out by the reparmetrization group (for genus $g = 0, 1$) and adding in the $6g - 6$ dimensional tangent space to Teichmüller space when $g > 1$. If B is represented for generic J then $\mathrm{Ind}(D_{u,J}) \geq 0$. Further, the adjunction formula implies that $g \leq 1 + \frac{1}{2}(B^2 - c_1(B))$. Therefore

$$0 \leq \operatorname{Ind}(D_{u,J}) \leq 2c_1(B) + 2 + B^2 - c_1(B) - 2 = c_1(B) + B^2 = d(B),$$

as claimed. Further, if $B^2 < 0$ then, because $d(B) = B^2 + c_1(B) \geq 0$, the adjunction formula $g(B) \leq 1 + \frac{12}{(}B^2 - c_1(B))$ for simple connected curves in class B (cf. [Mc91, Mc94i]) implies that $g(B) = 0$, and $B \in \mathcal{E}$ is the class of an exceptional sphere. If $B^2 = 0$ we again have equality in this formula, so that the curve is embedded and $g(B) = 0$ or 1, as claimed. $\square$

2.3. **Representing exceptional spheres.** With this preparation, we can now prove Proposition 1.2.5. By Gromov compactness, each class with $\mathrm{Gr}(A) \neq 0$ has some representative for every $J \in \mathcal{J}(\mathcal{S})$. In general this will be a nodal curve, i.e. a possibly disconnected image of a stable map. We denote these as Σ^A, reserving the notation C^A for a (smooth, often immersed) curve. When $J \in \mathcal{J}(\mathcal{S})$ a nodal curve Σ^A has some components in multiples of the classes S_i of the curves in $\mathcal{S}$ together with others that are possibly multiple covers of simple (i.e. somewhere injective) connected curves in some other classes B'_j. Thus the class A decomposes as

$$A = \sum_{i=1}^{s} \ell_i S_i + \sum_{j=1}^{k} n'_j B'_j, \tag{2.3.1}$$

where

- $\ell_i \geq 0$ and $n'_j > 0$ for all j,
- each class B'_j is represented by a simple J-holomorphic curve $C_j^{B'}$,
- the classes $B'_j, j = 1, \ldots, k$, are all distinct,
- $B'_j \neq S_i$ for any i, j.

Proof of Proposition 1.2.5. We saw in Lemma 2.2.3 that $\mathcal{J}_{SW}(\mathcal{S})$ is residual and path connected. We show below that when $J \in \mathcal{J}_{SW}(\mathcal{S})$ each $A \in \mathcal{E}$ is represented by a J-holomorphic embedded sphere. Then A can have no other representative with decomposition as in (2.3.1), for if it did A would have nonnegative intersection with each of its components, and hence with A itself. But $A^2 = -1$.

Suppose, to the contrary, that there is $A \in \mathcal{E}$ and $J \in \mathcal{J}_{SW}(\mathcal{S}, \omega, A)$ such that A has no embedded J-holomorphic representative. Since there is $J \in \mathcal{J}(M, \omega)$ such that *all* classes in $\mathcal{E}$ have embedded J-holomorphic representatives, there are only finitely many $E \in \mathcal{E}$ with $\omega(E) \leq \omega(A)$. Therefore we may choose $A_0 \in \mathcal{E}$ to be a class that satisfies the following conditions:

- $A_0 \cdot S_i \geq 0$ for all i;
- there is $J_0 \in \mathcal{J}_{SW}(\mathcal{S}, \omega, A_0)$ such that A_0 has no embedded J-holomorphic representative;
- $\omega(A_0)$ is minimal among all classes satisfying the previous conditions.

Fix such A_0, J_0. When $J \in \mathcal{J}_{reg}(M, \omega)$ is generic each class in $\mathcal{E}$ has a unique representative by an embedded J-holomorphic sphere. Because J_0 is the C^∞ limit of a sequence of elements $J_n \in \mathcal{J}_{reg}$, there is a J_0-holomorphic nodal curve $(\Sigma^{A_0})'$ in class A_0 whose components are all spheres, namely the Gromov limit of the unique J_n-holomorphic exceptional spheres $C_n^{A_0}$. Since $(\Sigma^{A_0})'$ is not embedded, the corresponding decomposition (2.3.1) is nontrivial, i.e. some $\ell_i \neq 0$ or $k > 1$. Further, because the integral of ω is positive on each component we have $\omega(B_j') < \omega(A_0)$ for all j. Thus none of the classes B_j' equal A_0. By definition of $\mathcal{J}_{SW}(\mathcal{S}, \omega, A_0)$ they have $c_1(B_j') > 0$ and $d(B_j') \geq 0$.

Since $A_0 \cdot A_0 < 0$ and $A_0 \cdot S_i \geq 0$ for all i, there must be some class $B_j' =: B''$ such that $A_0 \cdot B'' < 0$. Since $A_0 \cdot E \geq 0$ for all $E \in \mathcal{E} \smallsetminus \{A_0\}$, this class $B'' \notin \mathcal{E}$. Moreover, because the class B'' has a J-holomorphic representative for some $J \in \mathcal{J}(M)$ it can be represented by a symplectic submanifold; for details see Lemma 3.2.1 applied with $\mathcal{S} = \emptyset$. But then Lemma 2.1.4 implies that M is the blow up of a rational or ruled manifold. Because the class B'' is the J-holomorphic image of a sphere, we may now apply Lemma 2.1.3 parts (i) and (iii) to conclude that $\mathrm{Gr}(B'') \neq 0$. We now claim that B'' is reduced. This holds because, by the minimality of A_0, every class $E \in \mathcal{E}$ with energy $< \omega(A_0)$ is represented by a J_0-holomorphic embedded sphere and so has $E \cdot B'' \geq 0$. But then both classes A and B'' have embedded and connected J-holomorphic representatives for generic $J \in \mathcal{J}(M)$ and so have $A \cdot B'' \geq 0$, a contradiction. □

Remark 2.3.1. Of course, classes $E \in \mathcal{E}$ do degenerate, for example as $(E - E') + E'$ where $E' \in \mathcal{E}$. But such degenerations (a) happen for J in a set of codimension at least 2, and (b) have the property that the intersection of E with the class of the nonregular component(s) (in this case $E - E'$) is *negative*. The argument above shows the presence of nonregular components in classes S_i with $E \cdot S_i \geq 0$ does not affect the situation.

3. Geometric general position arguments

We next show how to maneuver the nodal curve $(\Sigma^A)'$ that represents A as in (2.3.1) so that its components are better behaved. After describing the patching procedures that we will use to resolve transverse intersections, we prove some basic general position lemmas and then Proposition 3.2.4, which is an important result about decompositions. As in §2, a curve is always assumed to have smooth domain. Otherwise it is called a stable or nodal curve.

3.1. Patching procedures.

Let $(\mathbb{C}^2, J_0, \omega_0)$ be Euclidean space with its usual complex structure J_0 and symplectic form ω_0.

Definition 3.1.1. *An immersed J-holomorphic curve C^B is* **orthogonal** *to $\mathcal{S}$ if there is $J \in \mathcal{J}(\mathcal{S})$ such that*

- *C^B is J-holomorphic,*
- *each intersection point $q \in C^B \cap \mathcal{S}$ has a neighborhood $\mathcal{N}_q$ such that there is a holomorphic symplectomorphism from $(\mathcal{N}_q, J, \omega)$ to a "square" neighborhood*

$$\mathcal{R}_\mu = \big\{(x, y) \mid |x|, |y| < \mu\big\} \subset (\mathbb{C}^2, J_0, \omega_0) \tag{3.1.1}$$

of $\{0\}$ such that the local J-holomorphic fibration $\mathcal{L}_{-k} \to \mathcal{S}$ is given by $(x, y) \mapsto x$, and $C^B \cap \mathcal{N}_q$ coincides with a fiber of this map.

Note that in this case there is only one branch of C^B through each point $q \in C^B \cap \mathcal{S}$. Note also that our hypotheses imply that the local Kähler structure (ω, J) near q (with $J \in \mathcal{J}(\mathcal{S})$) is flat. In the following we will resolve the singularities of orthogonally intersecting pairs of curves C^B, C^S by patching together the graphs of holomorphic sections σ_B, σ_S of the normal bundles to C^B and C^S. When C^B intersects C^S orthogonally, we can choose $J \in \mathcal{J}(\mathcal{S})$ so that both these bundles (not only that over C^S) have a complex structure, and so that the induced complex structure on a small neighborhood of $C^B \cup C^S$ can be extended to an ω-tame almost complex structure on M. We will consider sections that may have simple poles at the intersection point.

In the following we use the words **symplectic embedding** to mean a smooth embedding $f : \Sigma \to M$ such that the pullback form $f^*(\omega)$ does not vanish on the domain Σ, though otherwise the form $f^*(\omega)$ is arbitrary. Every J-holomorphic embedding is a symplectic embedding in this sense; moreover, any map f with this property is J-holomorphic for some ω-tame J.

It has long been understood that if two branches of a curve intersect positively and transversally, then one can get rid of this double point by a smooth surgery that replaces two discs by a cylinder (resolving the node): locally one replaces the graph of $xy = 0$ by $xy = \varepsilon$. This process allows one to build a symplectically embedded representative of a class B from an immersed curve with positive crossings. Our problem is more delicate, since if one of the initial curves lies in $\mathcal{S}$ we want the resulting glued curve to be disjoint from $\mathcal{S}$, or at most to intersect it orthogonally, so that it can be J-holomorphic for some $J \in \mathcal{J}(\mathcal{S})$. Lemma 3.1.3 describes the local patching procedure in its simplest form. We begin with a useful technical lemma.

Lemma 3.1.2. *Whenever $0 \le \delta_1 < \delta_0 \le r_0$ there is a symplectic embedding*

$$f : (r_0, 1) \times S^1 \to \mathcal{R}_1 \subset \mathbb{C}^2, \qquad (r, \theta) \mapsto (re^{i\theta}, \tfrac{\varepsilon(r)}{r}\, e^{-i\theta})$$

such that $\varepsilon(r) = \delta_1$ for r near 1 and $\varepsilon(r) = \delta_0$ for r near r_0.

Proof. We need $f^*(r_1\, dr_1 d\theta_1 + r_2\, dr_2 d\theta_2) > 0$. For this, we must choose $\varepsilon(r)$ so that

$$r - \frac{\varepsilon(r)}{r}\Big(\frac{\varepsilon(r)}{r}\Big)' \;=\; r - \frac{\varepsilon(r)}{r}\Big(-\frac{\varepsilon(r)}{r^2} + \frac{\varepsilon'(r)}{r}\Big) \;>\; 0$$

for $r_0 \le r \le 1$. This is always possible when $\delta_0 > \delta_1$ since we can then take $\varepsilon(r)$ to be a decreasing function of r. □

Lemma 3.1.3. *Suppose that the two J-holomorphic curves $C^S \subset \mathcal{S}$ and C^B intersect orthogonally at q, and that $J \in \mathcal{J}(\mathcal{S})$ is integrable (and fibered) near the point q on both curves. Let σ_S, σ_B be J-meromorphic sections of their normal bundles defined in some neighborhood $\mathcal{N}_q$ of q, each of which is either identically zero near q or is nonvanishing with a simple pole at q.*

(i) *If $\nu > 0$ is sufficiently small, the sections $\nu\sigma_S$ and $\nu\sigma_B$ can be patched together by a cylinder at q to form a (local) symplectically embedded curve C' that agrees with $\nu\sigma_S \cup \nu\sigma_B$ near the boundary $\partial\mathcal{N}_q$.*

(ii) *If σ_S has a pole at q, then we can choose ν so that C' is J'-holomorphic for some $J' \in \mathcal{J}(\mathcal{S})$. Further, if both sections have poles then C' is disjoint from $C^S \cup C^B$.*

Proof. By assumption we can identify $\mathcal{N}_q$ both symplectically and holomorphically with the region $\mathcal{R}_\mu := \{(x,y) \in \mathbb{C}^2 \mid |x|,|y| \le \mu\}$ for some $\mu > 0$, where $C^S = \{y = 0\}$, $C^B = \{x = 0\}$. Scaling up both x, y by a factor of $1/\mu$ we can assume that $\mu = 1$. For suitably small ν the sections σ_S, σ_B lie in $\mathcal{R}_1$ and have equations $xy \approx \delta_S, xy \approx \delta_B$, where the constants $\delta_B, \delta_S \ge 0$ are as small as we want. We divide into cases.

Case 1: *Neither section has a pole.* Then $\delta_S = \delta_B = 0$ and the equations are exact. We can then use Lemma 3.1.2 with $\delta_1 = 0, \delta_0 = \delta$ to interpolate between $xy = 0$ and $xy = \delta$ over $1/2 \le |x| \le 1$ and (interchanging the roles of x, y) between $xy = 0$ and $xy = \delta$ over $1/2 \le |y| \le 1$. If δ is sufficiently small, these curves are J'-holomorphic for J' arbitrarily $\mathcal{C}^1$-close to J. These two pieces can then be joined by the J-holomorphic curve $xy = \delta$.

Case 2: *Exactly one section has a pole.* Call the section that has a pole σ_S. Then we can arrange as above that it is given in $\mathcal{R}_1$ by an equation arbitrarily close to $xy = \delta_S$ for an arbitrarily small $\delta_S > 0$. In particular its graph over the annulus $1/2 \le |x| \le 1$ does not meet $C^S := \{y = 0\}$. Therefore we can smooth it out over $1/2 < |x| < 1$ by an arbitrarily $\mathcal{C}^1$-small perturbation so that it agrees with $xy = \delta_S$ for $|x|$ near $1/2$ without changing its intersection with a (very small) neighborhood of $\mathcal{S}$. Hence this piece of curve is J'-holomorphic for some $J' \in \mathcal{J}(\mathcal{S})$. Now interpolate between $xy = 0$ and $xy = \delta_S$ over $1/2 \le |y| \le 1$, and patch the two pieces by the J-holomorphic curve $xy = \delta_S$ as before. This proves (i) and (ii) in this case.

Case 3: *Both curves have poles.* The argument is similar. The graphs of each section should be first smoothed out so that they are given by the equations $xy = \delta_S$ and $xy = \delta_B$, and then the smaller constant should be increased to the larger one over an annulus for $|x|$ (or $|y|$) in the interval $[1/4, 1/2]$. The two pieces should then be joined by the J-holomorphic curve $xy = \max(\delta_S, \delta_B)$. □

Remark 3.1.4. Note that in this construction we can choose J' to be arbitrarily $\mathcal{C}^1$-close to J.

It will also be useful to consider meromorphic sections σ with poles of order $m > 1$. In this case they must be matched with the pushforward of a section by a branched cover.

Lemma 3.1.5. *Suppose that the two J-holomorphic curves $C^S \subset \mathcal{S}$ and C^B intersect orthogonally at q, and that $J \in \mathcal{J}(\mathcal{S})$ is integrable (and fibered) near the point q on both curves. Let σ_S be a J-meromorphic section as above with a pole of order m, and let σ_B be the graph of a multivalued meromorphic function over C^B that pulls back over the branched cover $z \mapsto x = z^m$ to a section with a simple pole. Then for sufficiently small ν the curves $\nu\sigma_S$ and $\nu\sigma_B$ can be patched together by a cylinder at q to form a (local) symplectically embedded curve C' that agrees with $\nu\sigma_S \cup \nu\sigma_B$ near the boundary $\partial\mathcal{N}_q$ and is disjoint from $C^S \cup C^B$.*

Proof. Note that the pushforward of the graph of the simple pole $y = \frac{1}{z}$ by $z \mapsto z^m =: x$ is the curve $y^m = \frac{1}{z^m} = \frac{1}{x}$; or equivalently $xy^m = 1$. But this can also be considered as the graph of the pole $x = \frac{c}{y^m}$ over C^S. Therefore these two curves can be patched together as in Lemma 3.1.3. The result for arbitrary sections σ_S, σ_B of the given form follows as in Lemma 3.1.3. $\square$



FIGURE 3.1.1. As in Example 3.1.6 we glue C^S to a triple cover of C^E by matching two poles of order 3 and one simple pole of σ_S with corresponding parts of the multisection σ_E as labelled. The gluing is indicated by dotted lines, the three relevant parts of C^E are shown in black, while the two other branches of σ_B near the simple pole of σ_S are shown by thinner lines.

The next example illustrates the use of Lemma 3.1.5.[4]

Example 3.1.6. Suppose that $\mathcal{S}$ consists of a single sphere in class S with $S \cdot S = -7$ and that E is the class of an exceptional divisor with $E \cdot S = 3$. Then $A := S + 3E$ has $A^2 = 2, c_1(A) = -2$ so that $d(A) = 0$ and $g(A) = 3$. Further $A \cdot E = 0$ and $A \cdot S = 2$. Hence we should be able to represent A by an embedded genus 3 curve that meets $\mathcal{S}$ in two points. We will assume (as we may by Lemma 3.2.1 below) that C^E meets C^S in three distinct points, and will parametrize C^E so that these points are $0, 1, \infty$. Consider the map $f : S^2 \to C^E, z \mapsto z^3$. The pullback $f^*(\mathcal{L}_{-1})$ of the normal bundle to C^E has Chern class -3 and so has a section σ with simple poles at $0, 1, \infty$. Let σ_S be a nonvanishing meromorphic section of $\mathcal{L}^S_{-7}$ with poles of order three at both $0, \infty$ and a simple pole at 1. By Lemma 3.1.5 we can patch σ_S to the pushforward of σ at $0, 1, \infty$ to obtain a closed curve C in class $S + 3E$. Note that, as required, C intersects C^S in two points, one for each of the other two points in $f^{-1}(1)$: see Figure 3.1.1. More precisely, these points correspond to $\sigma(\gamma(1)), \sigma(\gamma^2(1))$, where

$$\gamma : S^2 \to S^2, \quad z \mapsto e^{\frac{2\pi i}{3}} z,$$

---

[4] I want to thank Emmanuel Opshtein for pointing out a problem with my first attempt to patch such curves together.

so that $f\circ\gamma = f$. These values $\sigma(\gamma(1)), \sigma(\gamma^2(1))$ are very small: hence they lie in the disc in C^S that is cut out when we do the gluing at the point $1 \in C^E$. Therefore the glued curve C has no self intersections near C^S. As another check that this is correct, note that if they did not lie in this disc then the pushforward $f_*(\sigma)$ would have a self-intersection near $1 \in C^E$, which is not the case. In fact, any two distinct meromorphic sections of $\mathcal{L}_{-3}$ meet in three points. But the sections σ and $\sigma^\gamma := \sigma\circ\gamma$ of $\mathcal{L}_{-3}$ have two of their intersection points at the poles over $0, \infty$ and so have only one other intersection point. Hence there is a unique solution $w_1 \in \mathbb{C}$ to the equation $\sigma(w) = \sigma^\gamma(w)$. But then the unique solution to the equation $\sigma(w) = \sigma^{\gamma^2}(w)$ is $w_2 = \gamma w_1$, and the unique solution to the equation $\sigma^\gamma(w) = \sigma^{\gamma^2}(w)$ is $w_3 = \gamma^{-1}w_1$. These three intersection points correspond to a single self-intersection of the pushforward section $f_*(\sigma)$. Hence the glued curve C has precisely one self-intersection. Moreover, because it was formed by gluing together two copies of $S^2 -$ *three discs*, C has genus 2. Hence it can be smoothed to an embedded curve of genus 3. This is consistent with the fact that $g(A) = 3$.

Another possibility would be to use the branched covering map $f : T^2 \to S^2 = C^E$ of degree 3 with exactly three totally ramified branch points that is provided by Lemma 3.1.7 below, take a section ρ of $f^*(\mathcal{L}^E_{-1})$ that has a simple pole at each of the branch points, and patch its pushforward $f_*(\sigma)$ with a section σ of $\mathcal{L}^S_{-7}$ that has three poles of order three and two zeros.



FIGURE 3.1.2. (I) shows the domain Σ of a two-fold branched cover of the sphere with 6 branch points; the map quotients out by a half turn through the line through the branch points. (II) shows a schematic picture for the domain of a branched cover of S^2 with even degree $m = 4$. To get m branch points, one needs $\frac{m}{2} - 1$ "beads" (three are shown), Σ is the boundary of a tubular neighborhood of this one-complex, and the map quotients out by a rotation through angle $2\pi/m$ about the axis of symmetry.

The following lemmas will be useful later.

Lemma 3.1.7. *For each $m \geq 2$, there is a degree m branched covering map $f : \Sigma \to S^2$ of S^2 by a closed Riemann surface Σ with precisely m branch points, each of which is totally ramified i.e. has local model $z \mapsto z^m$.*

Proof. When $m = 2b$ is even, one can easily construct such a cover as in Figure 3.1.2. In general, this is a classical result due to Hurwitz [H91]. One just needs to construct a covering of $S^2 \smallsetminus m$ *points* of degree m whose monodromy round each deleted point is the cyclic permutation $(12\cdots m)$; see [EKS84, Lemma 2.1]. □

Lemma 3.1.8. *Suppose that C_1, C_2 are orthogonally (and hence positively) intersecting embedded spheres, and let σ_1, σ_2 be small, (not necessarily holomorphic) sections of their normal bundles that are normalized near the intersection points as in Lemma 3.1.3 either to be constant or to have a simple pole. If $C_1 \cdot C_2 = 1$ and the unique intersection point is resolved, then the resulting embedded curve is a sphere. If $C_1 \cdot C_2 > 1$ and all intersection points are resolved then the resulting embedded curve has genus > 0.*

Proof. The topological effect of the patching process is the same whether the sections have a simple pole or are constant: it removes a small disc from each curve through the intersection point and replaces this by a cylinder $S^1 \times [0,1]$. Thus the embedded curve is obtained from the disjoint union of two 2-spheres by k disjoint positively oriented surgeries of index 1 (or equivalently by taking the connected sum at k different points), where k is the number of points resolved. The first such surgery makes a single sphere, and each subsequent surgery adds one to the genus. □

3.2. **General position arguments.**

The following general position argument is well known: we include it for completeness.

Lemma 3.2.1. *Let $B = \sum_{j=1}^{t} m_j B_j \in H_2(M)$ be such that*

- *$B_j \neq B_k$ for each $j \neq k$;*
- *$B_j \cdot S_i \geq 0$ for all i, j;*
- *$m_j \geq 1$and $= 1$ if $B_j^2 < 0$;*
- *for some $J_0 \in \mathcal{J}(\mathcal{S})$, each B_j can be represented by a simple J_0-holomorphic curve.*

Then B can also be represented by a (possibly disconnected) embedded curve that is orthogonal to $\mathcal{S}$ and J-holomorphic for some $J \in \mathcal{J}(\mathcal{S})$.

Proof. First, consider the case when $t = m_1 = 1$, so that there is a simple J_0-holomorphic curve C^B. It has at most a finite number of singular points $q_i = u(z_i)$. Suppose first that none lie on $\mathcal{S}$. At each of these it is possible to perturb C^B locally to an immersed J_0-holomorphic curve by [Mc94i, Theorem 4.1.1], and then patch this new piece of curve to the rest of C^B by the technique of [Mc91, Lemma 4.3], to obtain a positively immersed symplectic curve. This curve $(C^B)'$ is J_0-holomorphic except close to $C^B \cap Shell$, where $Shell$ is the union of spherical shells $Shell(q) := B_{r_1}(q) \smallsetminus B_{r_2}(q)$ centered at the finite number of singular points q. Thus we can make it J-holomorphic for some J near J_0 that equals J_0 away from $(C^B)' \cap Shell$. Hence even if some singular point q is in C^S we can assume $J \in \mathcal{J}(C^S)$. Indeed, if there is a singularity $q \in C^S$ we can avoid the analytic argument in [Mc94i], since J_0 is integrable and fibered near q. Hence the curve C^B is the graph of some holomorphic section of the normal bundle $\mathcal{L}^S_{-k}$ to $\mathcal{S}$ with a zero of order > 1 at q, and it suffices to perturb this to a J_0-holomorphic section with transverse zeros and then patch the result to the original curve by a patch that does not intersect the zero set of $\mathcal{L}^S_{-k}$.

Then $(C^B)'$ is immersed, and can be homotoped (keeping it symplectic) so that it has at most transverse double points that are disjoint from its intersections with the spheres C_i^S in $\mathcal{S}$. Then we deform $(C^B)'$ so that it is vertical near its intersections p with each C_i^S, in the sense that it coincides with the fiber of the normal bundle $\mathcal{L}^S_{-k_i}$ at p. This maneuver is carried out in detail in [Mc94ii, Lemma 3.11]. (We also have to be careful to keep it J_0-holomorphic near C_i^S; but this is not difficult.) Then $(C^B)'$ meets each component C_i^S of $\mathcal{S}$ orthogonally in distinct points. Moreover, by resolving all its double points (which lie away from C_i^S) as in Lemma 3.1.3, we can assume that $(C^B)'$ is embedded and still J-holomorphic for some $J \in \mathcal{J}(\mathcal{S})$. This completes the proof when B has a simple representative C^B.

Now suppose that $B = mB_0$ where B_0 has a simple representative C^{B_0}. By the above we can suppose that C^{B_0} is embedded, orthogonal to $\mathcal{S}$ and J-holomorphic for some $J \in \mathcal{J}(\mathcal{S})$. Then for suitable $J \in \mathcal{J}(\mathcal{S})$ a neighborhood $\mathcal{N}(C^{B_0}, J)$ of C^{B_0} can be identified with a neighborhood of the zero section in a complex line bundle over C^{B_0} with nonegative Chern class. Hence we may represent the class mB_0 by the union of m generic J-holomorphic sections of this bundle that intersect transversally. If $B_0^2 > 0$ each pair of these sections intersect, and by choosing generic sections we can assume that the intersection points do not lie on $\mathcal{S}$. Hence after resolving these intersections as before, we get an embedded (possibly disconnected) representative of mB_0 that we finally perturb to be orthogonal to $\mathcal{S}$.

This completes the proof when $t = 1$. If $t > 1$ we first resolve all singularities, so that each simple curve C^{B_j} is immersed and meets the other C^{B_k} as well as $\mathcal{S}$ transversally and positively in double points. Next, we perturb these double points to be orthogonal. We then replace the components with $m_j > 1$ by suitable nearby embedded curves in class m_jB_j. Finally, we patch all double points to get an embedded curve in class B. □

The following extension of this lemma to multiply covered exceptional classes will also be useful.

Lemma 3.2.2. *Let $J \in \mathcal{J}(\mathcal{S})$ and suppose that C^B and C^E are transversally intersecting embedded J-holomorphic curves orthogonal to $\mathcal{S}$, where $B \cdot E = k \geq 1$ and $E \in \mathcal{E}$. Then, for each $1 \leq m \leq k$, there is $J' \in \mathcal{J}(\mathcal{S})$ that equals J on TC^E and an embedded J'-holomorphic curve C in class $B + mE$ that meets C^E orthogonally in exactly $k - m$ points.*

Proof. First isotop C^B so that it meets C^E transversally at points in $C^E \smallsetminus \mathcal{S}$, and then change J near C^E (but not on the tangent bundle TC^E) to an element $J' \in \mathcal{J}(\mathcal{S})$ that is integrable and fibered near C^E. Thus we can identify a neighborhood $\mathcal{N}(C^E)$ of C^E in M with a neighborhood of the zero section in the holomorphic normal line bundle $\mathcal{L}^E_{-1}$ to C^E. Since the intersection points $C^B \cap C^E$ do not lie on $\mathcal{S}$ we can also assume that J' is normalized near these intersection points to have the product form $\mathcal{R}_\mu$ as in Definition 3.1.1, and then manipulate C^B further so that it is orthogonal to C^E, intersecting it at the points $q_1, \dots, q_k$. Since $|C^B \cap C^E| = k \geq 1$ we may choose a (small) nonvanishing section σ of $\mathcal{L}^E_{-1}$ with a simple pole at q_k, and then use Lemma 3.1.3 to patch C^B to the graph $graph(\sigma)$ of this section. Since $\sigma(q_i) \neq 0$ when $i < k$, by using very small patches at $\sigma(q_1), \dots, \sigma(q_{k-1})$ we may do this without changing J' on C^E or changing C^B very near the intersection points $q_1, \dots, q_{k-1}$. Therefore, the resulting embedded curve lies in class $B + E$ and intersects C^E orthogonally at the points $q_1, \dots, q_{k-1}$. Repeating this process $m - 1$ more times (with smaller sections σ each time) we finally construct the desired curve in class $B + mE$. □

Remark 3.2.3. To prove Proposition 1.2.9 we will need to carry out geometric procedures similar to those in Lemmas 3.1.3, 3.2.1 and 3.2.2 above for generic smooth 1-parameter families of nodal curves whose components are immersed and lie in given homology classes. Since the choices at each stage are contractible, there is no problem with doing this.

We now explain how to "clean up" a nodal representative of the class A. As explained earlier, when $J \in \mathcal{J}(\mathcal{S})$ for each such stable map $(\Sigma^A)'$ there is a corresponding decomposition of the class A:

$$A = \sum_{i=1}^{s} \ell_i S_i + \sum_{j=1}^{k} n'_j B'_j, \tag{3.2.1}$$

where $\ell_i \geq 0$, $n'_j > 0$, each class B'_j is represented by a simple J-holomorphic curve $(C^B_j)'$, and $B'_j \neq S_i$ for any i, j. Note that we require J' to be a generic element in $\mathcal{J}(\mathcal{S})$ in order to control the class B in (3.2.2).

Proposition 3.2.4. *Suppose that $b_2^+(M) = 1$ and that the class A satisfies the conditions of Theorem 1.2.1. Let $(\Sigma^A)'$ be a J'-holomorphic nodal curve in class A with corresponding decomposition (3.2.1). Then If $J' \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$, there is $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ such that the class A is represented by a connected J-holomorphic nodal curve Σ^A with components of the following types, where the ℓ_i are as in (2.3.1):*

- *an ℓ_i-fold cover of S_i for $i = 1, \dots, s$;*
- *m_j-fold coverings of exceptional spheres C^E_j in classes $E_1, \dots, E_e$ for $e \geq 0$ that are mutually disjoint and orthogonal to $\mathcal{S}$;*
- *an embedded curve C^B in the class B, that is*
  - *connected, unless $B^2 = 0$, and $B = mB_0$ where $m > 1$, in which case it consists of m parallel copies of an embedded sphere or torus in class B_0;*
  - *orthogonal to $\mathcal{S}$ and disjoint from the spheres C^E_i.*

In particular

$$A = \sum_{i=1}^{s} \ell_i S_i + \sum_{j=1}^{e} m_j E_j + B, \tag{3.2.2}$$

where B is reduced, $B^2 \geq 0$, $d(B) \geq 0$, $m_j > 0$, $E_j \in \mathcal{E}$ and $E_j \cdot E_{j'} = E_j \cdot B = 0$ for all $j \neq j'$ with $j, j' \in \{1, \dots, e\}$.

Proof. Suppose first that $J' \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$, and consider the components of the original nodal curve $(\Sigma^A)'$ that are not in $\mathcal{S}$. These are n'_j-fold covers of simple curves in classes B'_j, where $d(B'_j) \geq 0$ for all i by Lemma 2.2.4 (ii) and our hypothesis on J'. By Lemma 3.2.1, we can assume that the curves in classes B'_j are embedded, that they meet transversally, and that they are orthogonal to $\mathcal{S}$. We may further assume that $n'_j = 1$ except possibly if $(B'_j)^2 \leq 0$. But if $(B'_j)^2 = 0$, then the curve in class B'_j has trivial normal bundle so that we can replace its n'_j-fold cover by n'_j parallel (and disjoint) copies of the B'_j-curve. By Lemma **??** (ii), the classes B'_j with $(B'_j)^2 < 0$ lie in $\mathcal{E}$. Relabel the classes B'_j that lie in $\mathcal{E}$ as $E_j, j = 1, \dots, e'$, and call the remaining (distinct) classes $B'_j, j = 1, \dots, k'$. These other classes all have $(B'_j)^2 \geq 0$ and therefore by Lemma 2.1.1 if $k' > 1$ these classes must all intersect. Hence by Lemma 3.2.1, in this case we may amalgamate all these curves into one embedded curve $C^{B'}$ in a class B' with $(B')^2 > 0$. Thus in all cases, we can assume that $k' = 1$, and that the class B' is represented by an embedded curve that is connected except possibly if $(B')^2 = 0$. Hence the decomposition is:

$$A = \sum_{i=1}^{s} \ell_i S_i + \sum_{j=1}^{e'} m'_j E'_j + B'. \tag{3.2.3}$$

If $B' \cdot E'_j = 0$ and $E'_i \cdot E'_j = 0$ for all i, j, then the nodal curve Σ^A is in the required form. Therefore we take $E_j := E'_j$ and $B := B'$, and it remains to check that we can assume Σ^A is holomorphic for some $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$. But the exceptional spheres C^E_j are automatically regular, and the B curve is embedded and has $d(B) \geq 0$ by construction. Hence this follows by Lemma 2.2.4 (i).

We now complete the proof in the case when $J' \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ by showing how to arrange that $B' \cdot E'_j = 0$ and $E'_i \cdot E'_j = 0$ for all i, j. If $E'_i \cdot E'_j > 0$ for some $i \neq j$ then we can reduce the number e' of exceptional classes in the decomposition (3.2.3) as follows. Suppose without loss of generality that $m'_i \leq m'_j$. By Lemma 3.2.2 we may form an embedded curve C in class $[C] = E'_i + E'_j$ that is J''-holomorphic for some $J'' \in \mathcal{J}(\mathcal{S})$ for which $C^{E'}_j$ is also holomorphic. Then $C^2 \geq 0$ and so by Lemma 2.1.1 if $B' \neq 0$ then either $[C] \cdot B' > 0$ or $[C]^2 = 0$ and B' and $[C]$ are multiples of one another. In the former case, we can use the method of Lemma 3.2.1 to make an embedded curve in class $B'' := B' + m'_i[C]$ that is J''-holomorphic for some possibly different $J'' \in \mathcal{J}(\mathcal{S})$ for which all the curves $C^{E'}_j$ except perhaps $C^{E'}_i$ are holomorphic. Since the class $m'_i E'_i$ is now incorporated into the B'' curve, this reduces e' by at least one. (It is reduced by two if $m'_i = m'_j$.) In the latter case, we can also reduce e' by incorporating a curve in class $m'_i[C]$ into the B'-curve; indeed the condition $[C]^2 = 0$ implies that $E'_i \cdot E'_j = 1$ so that C is an embedded sphere, and since $[C] \cdot B' = 0$, the B'-curve must consist of some number of parallel copies of C to which we can add more copies. Finally, if $B' = 0$ then we can use $m'_i[C]$ as the new class B'', which again reduces e'. By repeating this process, we can therefore reduce to the case when all the exceptional spheres $C^{E'}_j$ are disjoint. Therefore we now define $E_j := E'_j$.

The final step is to use Lemma 3.2.2 to reduce the number

$$\kappa := \sum_j B' \cdot m_j E_j.$$

If $\kappa = 0$ there is nothing to prove. So suppose that $B' \cdot E_{j_0} = k > 0$. If $k \le m_{j_0}$, we can create an embedded curve C in class $B'' := B' + kE_{j_0}$ that is disjoint from E_{j_0} and so replace the components in classes $B', m_{j_0}E_{j_0}$ by ones in classes $B'', (m_{j_0} - k)E_{j_0}$. Since $E_{j_0} \cdot E_j = 0$ for all $j \ne j_0$ the intersections of the B'' curve with the classes E_j, for $j \ne j_0$ are unchanged. Hence this reduces κ by k. On the other hand, if $k \ge m_{j_0}$ we form a curve B'' in the class $B' + m_{j_0}E_{j_0}$ and remove E_{j_0} from the list of exceptional classes. This reduces κ by m_{j_0}. Continuing in this way, we eventually arrive at a situation in which $\kappa = 0$. This completes the proof. □

Lemma 3.2.5. *When the decomposition* (3.2.2) *is obtained from* (3.2.1) *as above,* $d(B) \ge \sum_{j=1}^k \max\big(0, d(n'_j B'_j)\big)$.

Proof. The class B for the decomposition (3.2.1) is obtained in stages, each step replacing components that lie in classes D, D' by a single component in class $D + D'$. These steps are of two kinds:

- either $D = mE$ and $D' = mE'$ where $E, E' \in \mathcal{E}$ have $E \cdot E' \ne 0$; or
- $D^2 \ge 0, (D')^2 \ge 0$ and $D \cdot D' > 0$.

For each $D \in H_2(M;\mathbb{Z})$, define $k(D) := \max(0, \frac{1}{2}d(D))$. One can interpret this as the number of generic points that a generic J-holomorphic curve in class D goes through. For example, when $D = mE$ we have $d(mE) = -m^2 + m \le 0$, so that $k(mE) = 0$, which corresponds to the fact that exceptional spheres are rigid. On the other hand if $D^2 \ge 0$ and $m, m' > 0$ then $d(mD) + d(m'D) \le d\big((m + m')D\big)$, which illustrates the fact that d (and hence k) increases as nonnegative curves are combined.

For each decomposition $\mathcal{D}$ of A as in (3.2.1), let us define $k(\mathcal{D}) := \sum_j k(n'_j B'_j)$. Then we claim that if $\mathcal{D}'$ is obtained from $\mathcal{D}$ by one of the two kinds of steps above, we have $k(\mathcal{D}') \ge k(\mathcal{D})$. This is clear for steps of the second kind since the intersections of the classes D, D' with all other components are nonnegative. To check it for a step of the first kind, note that if $E = E_1, E' = E_2$ then we choose $m := \min(m_1, m_2)$. Since the classes E_i do not contribute to $k(\mathcal{D})$, we have $k(\mathcal{D}') - k(\mathcal{D}) = \max(0, K)$ where

$$K := k\big(m(E_1 + E_2)\big) + \frac{m}{2}(E_1 + E_2) \cdot \big(A - m(E_1 + E_2)\big).$$

But $(E_1 + E_2)^2 \ge 2E_1 \cdot E_2 - 2 \ge 0$ and $c_1(E_1 + E_2) = 2$ so that $k\big(m(E_1 + E_2)\big) > 0$. Further $(E_1 + E_2) \cdot E_i \ge 0$ since $E_1 \cdot E_2 > 0$. Hence, because $m \le m_i$ for $i = 1, 2$, the second term is nonnegative also. This completes the proof. □

Definition 3.2.6. *We say that a stable map* Σ^A *is* **$\mathcal{S}$-normalized** *if it satisfies the conclusions of Proposition 3.2.4. In particular, its homology class A decomposes as in* (3.2.2).

Remark 3.2.7. Suppose that $\ell_i \ge 0$ are multiplicities such that $A' := A - \sum_{i=1}^s \ell_i S_i$ has $\mathrm{Gr}(A') \ne 0$. Then there is no easy way to guarantee that A is represented by a nodal curve $(\Sigma^A)'$ with precisely these multiplicities along $\mathcal{S}$, since every J-holomorphic representative of the class A' with $J \in \mathcal{J}_{reg}(\mathcal{S})$ might have some components in $\mathcal{S}$; cf. Corollary 4.1.3. What we are claiming above is that if A has such a representative, then we can further arrange that the components of A' are as they would be for generic $J \notin \mathcal{J}(\mathcal{S})$, i.e. there is a main embedded component C^B in a reduced class together with some multiply covered and disjoint exceptional spheres, with the added bonus that all these components are orthogonal to $\mathcal{S}$. The proof is purely geometric. In particular, it does not argue that because $\mathrm{Gr}(A') \neq 0$ the required decomposition follows by considering a generic J-holomorphic curve through $d'(A')$ points, as we did earlier (cf. the discussion around (2.1.1)). In fact because we would have $J \in \mathcal{J}(\mathcal{S})$, such an argument would be circular: we are trying to figure out what such a representative would look like for generic $J \in \mathcal{J}_{reg}(\mathcal{S})$, and so cannot assume anything about this during the course of the proof.

4. PROOFS OF THE MAIN RESULTS

4.1. **Proof of Theorem 1.2.1.** We convert the local geometric procedures described in §3.1 into global gluing results. We assume throughout the class A satisfies the conditions of Theorem 1.2.1. As a warmup we begin with some special cases.

Lemma 4.1.1. *Let $\mathcal{S}$ consist of a single curve C^S of self-intersection $-k \leq -2$ and $J_0 \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$, and suppose that in the decomposition* (3.2.2) *of A in Proposition 3.2.4 all $m_i = 0$. Then, there is $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ such that A has an embedded J-holomorphic representative that meets C^S orthogonally in $A \cdot S$ distinct points.*

Proof. By assumption $A = \ell S + B$ where $d(B) \geq 0$. We argue by induction on ℓ. If $\ell = 0$ there is nothing to prove. If $\ell > 0$, then $B \cdot S = A \cdot S - \ell S \cdot S \geq k\ell$, and the nodal curve Σ^A in Proposition 3.2.4 contains an embedded curve C^B that meets C^S in $\geq k\ell$ distinct points. Choose k of these points $p_1, \ldots, p_k$, and a nonvanishing meromorphic section σ of the normal bundle $\mathcal{L}^S_{-k}$ to C^S with k simple poles at $p_1, \ldots, p_k$ on C^S. Then for sufficiently small ν the graph of $\nu\sigma$ can be patched to the curve C^B (the graph of the zero section) near each $p_1, \ldots, p_k$ by Lemma 3.1.3. We obtain an immersed curve C' in class $B' = B + S$ that agrees with $C^B \cup \mathrm{graph}(\sigma)$ away from the points $p_1, \ldots, p_k$. Since σ does not vanish, the intersections of C' with C^S occur at the points $C' \cap C^S = C^B \cap C^S \smallsetminus \{p_1, \ldots, p_k\}$, and so are orthogonal by assumption. Using Lemma 3.2.1, we now resolve the new double points of C' (corresponding to intersections of C^B with $\mathrm{graph}(\sigma)$), to get an embedded curve $(C^B)'$ in class $B' = S + B$ that is orthogonal to C^S, and J-holomorphic for some $J \in \mathcal{J}(\mathcal{S})$. Then

$$d(B') = d(S) + d(B) + 2S \cdot B \geq 2 - 2k + 2k\ell > 0.$$

Therefore, by Lemma 2.2.4 (i) there is $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ such that the embedded curve $(C^B)'$ is J-holomorphic. Since $A = (\ell - 1)S + B'$, this completes the inductive step. □

Lemma 4.1.2. *Let $\mathcal{S}$ consist of a single curve C^S of self-intersection $-k \leq -2$ and $J_0 \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$, and suppose that $A = S + mE$, where $m \geq 1$ and E is represented by an exceptional sphere C^E. Then, there is $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ such that A has an* *embedded smooth J-holomorphic representative C^A that is transverse to both C^S and C^E.*

Proof. Let $a := E{\cdot}S$. Then $a > 0$, since A is reduced by condition (iii) in Theorem 1.2.1. We have $A = mE + S$ where by hypothesis

$$A \cdot S = ma - k \geq 0, \qquad A \cdot E = -m + a \geq 0,$$

the latter holding because $A \neq E$. Further, $a > 1$ since otherwise we have $k \leq m \leq 1$ which is impossible when $k \geq 2$.

By Lemma 3.2.1 we may assume that the embedded curve C^E meets C^S orthogonally in a points $q_1, \dots, q_a$, and choose $J \in \mathcal{J}(\mathcal{S}) \cap \mathcal{N}(J_0)$ so that a neighborhood of C^E is biholomorphic to a holomorphic line bundle $\mathcal{L}^E_{-1} \to C^E$. We use the method explained in Example 3.1.5. If $m = 1$ then $k \leq a$ so that there is a meromorphic section of $\mathcal{L}^S_{-k}$ with simple poles at $q_1, \dots, q_k$ that we can patch to a suitable meromorphic section of $\mathcal{L}^E_{-1}$ to obtain the desired curve C^A.

So suppose that $m > 1$. By Lemma 3.1.7 there is a branched cover $f : \Sigma \to S^2$ of S^2 of degree m by a closed Riemann surface Σ with precisely m branch points, each of which is totally ramified. Simce $m \leq a$ we may choose f so that its branch points map to the intersection points $C^S \cap C^E = \{q_1, \dots, q_m\}$. Let ρ be a nonvanishing meromorphic section of $f^*(\mathcal{L}^E_{-1})$ with simple poles at the branch points $f^{-1}(q_1), \dots f^{-1}(q_m)$. Then, as in Lemma 3.1.5, the pushforward multisection $f_*(\rho)$ may be matched to poles of order m at each of $q_1, \dots, q_m$ and it also intersects C^S transversally at $m(a-m)$ other points, m near each of the remaining intersection points $q_{m+1}, \dots, q_a$. Since $k \leq ma$ we may choose a meromorphic section σ of $\mathcal{L}^S_{-k}$ with poles of order m at each of $q_1, \dots, q_m$ and other poles at the intersection points of $f_*(\rho)$ with C^S as needed. (If $k < m^2$ the section σ will have some zeros as well.) Since $k \leq ma$, it is possible to place all the needed poles of σ. We now patch these sections as in Lemma 3.1.5 to form a symplectically immersed curve in class $A = S + mE$ that is holomorphic near $C^S \cup C^E$, and whose only singularities are transverse positive double points. We then resolve intersections using Lemma 3.2.1 to obtain an embedded representative for A that is J-holomorphic for some $J \in \mathcal{J}(\mathcal{S})$, that we can perturb to lie in $\mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ by Lemma 2.2.4 (i). □

With these preliminaries in place, we now explain the proof of Theorem 1.2.1 when $\mathcal{S}$ has just one component.

Proof of Theorem 1.2.1 when $\mathcal{S}$ has one component C^S: Since A is reduced and by hypothesis $\mathrm{Gr}(A) \neq 0$, for each $J \in \mathcal{J}(\mathcal{S})$ the class A has some connected J-holomorphic representative. Thus there is a corresponding decomposition (2.3.1) and nodal curve $(\Sigma^A)'$ as in the hypothesis of Proposition 3.2.4. In particular, the component in $\mathcal{S}$ represents the class $\ell' S$ for some $\ell' \geq 0$. If $\ell' = 0$, then because $(\Sigma^A)'$ is connected, we may convert it into a single embedded curve C^A by Lemma 3.2.1. We now suppose inductively that Theorem 1.2.1 holds if $(\Sigma^A)'$ has $0 \leq \ell' < \ell$, and consider the case $\ell' = \ell$. Applying Proposition 3.2.4, we may therefore assume that A has an $\mathcal{S}$-normalized representative $\Sigma^A = (\Sigma^A)_0$ in the sense of Definition 3.2.6, and will produce a representative of the form $(\Sigma^A)' = (\Sigma^A)'_1$ for some different decomposition with $\ell' < \ell$. The desired curve C^A then exists by the inductive hypothesis.

Lemma 4.1.1 considers the case when the $\mathcal{S}$-normalized nodal curve Σ^A involves no exceptional spheres. The remaining case is when there are some multiply covered exceptional spheres. For this, we must generalize Lemma 4.1.2 to the case when Σ^A has

- some (possibly multiply covered) disjoint exceptional spheres in classes m_jE_j, $j = 1, \dots, e$, with $m_j \geq 1$,
- a possibly disconnected embedded curve C^B that is disjoint from the E_j-curves, as well as
- some copies of C^S with total multiplicity ℓ.

Then $A = \sum_{j=1}^e m_jE_j + B + \ell S$, and $a_j := E_j \cdot S \geq 1$ because Σ^A is connected. Hence we have

$$A \cdot S = \sum m_ja_j + h - \ell k \geq 0, \quad A \cdot E_j = -m_j + \ell a_j \geq 0. \tag{4.1.1}$$

As explained above, we will argue by induction over ℓ. If

$$h := B \cdot S \geq k,$$

then we can use Lemma 4.1.1 to resolve k of the intersections of the curves C^B with C^S, thus reducing ℓ by one. Therefore we may assume that $h < k$.

Further if $e + h \geq k$ then we may construct an immersed curve in class $B' := S + \sum_{j=1}^{k-h} E_j + B$ by choosing for each $j = 1, \dots, k-h$ a nonvanishing meromorphic section ρ_j of $\mathcal{L}_{-1}^{E_j}$ with pole at q_j and a nonvanishing meromorphic section σ of $\mathcal{L}_{-k}^S$ with poles at the points $q_1, \dots, q_{k-h}$ as well as at $C^B \cap C^S$, and then patching σ to the union of C^B with the graphs of the ρ_j . The resulting curve C' meets C^S transversally near the other points of intersection of C_j^E with C^S. Further since $B^2, k-h \geq 0$,

$$(B')^2 = B^2 - (k-h) - k + \sum_{j=1}^{k-h} 2a_j + 2h \geq -2k + 3h + 2(k-h) = h \geq 0,$$

since each $a_j \geq 1$. Further it meets C_j^E positively for any $j \leq k-h$ with $a_j > 1$. Hence we may incorporate these curves C_j^E into C^B as in Lemma 3.2.1, and proceed with reduced ℓ and e.

The remaining case is when $e + h < k$. We first claim there are integers $1 \leq m_j' \leq \min(m_j, a_j)$ for $j = 1, \dots, e$ so that

$$k \leq \sum_{j=1}^e m_j'a_j + h. \tag{4.1.2}$$

To see this, assume without loss of generality that $a_1 \geq a_2 \geq \dots \geq 1$. If $m_j' = 1$ for all j, then because $e + h < k$ we have $\sum_{j=1}^e m_j' \leq k$. We now increase the m_j' for $j = 1$ in steps of size 1 to its maximum value $\min(m_j, a_j)$, then repeat this for $j = 2$, then $j = 3$ and so on. Since the sum $\sum_{j=1}^e m_j'$ increases by 1 each time, we either attain equality

$\sum_{j=1}^e m'_j = k$ at some stage, in which case $k \le \sum_{j=1}^e m'_j a_j + h$ since $a_j \ge 1$ and $h \ge 0$, or we have $\sum_{j=1}^e \min(m_j, a_j) < k$. In that case, using $m_j \le \ell a_j$ and $m'_j = \min(m_j, a_j)$, we have

$$\begin{aligned} k \;&\le\; \tfrac{1}{\ell}\Big(\sum_{m_j \le a_j} m_j a_j + \sum_{m_j > a_j} m_j a_j\Big) + \tfrac{h}{\ell} \\ &\le\; \tfrac{1}{\ell}\Big(\sum_{m_j \le a_j} m'_j a_j\Big) + \tfrac{1}{\ell}\Big(\sum_{m_j > a_j} \ell a_j a_j\Big) + h \\ &\le\; \sum_{m_j \le a_j} m'_j a_j + \sum_{m_j > a_j} m'_j a_j + h \\ &=\; \sum_{j=1}^{e} m'_j a_j + h, \end{aligned} \tag{4.1.3}$$

as required. This proves the claim.

Let $C_j^E \cap C^S = \{q_{j1}, \ldots, q_{j\,a_j}\}$. For each j choose a branched covering map $f_j : \Sigma_j \to C_j^E$ of degree m'_j with exactly m'_j totally ramified branch points with images $q_1, \ldots, q_{m'_j}$, as in Lemma 3.1.7. (If $m'_j = 1$, take f_j to be the identity map with "branch point" at q_1.) Choose a nonvanishing meromorphic section ρ_j of $f_j^*(\mathcal{L}_{-1}^{E_j})$ with simple poles at each branch point, and consider its pushforward $(f_j)_*(\rho_j)$. As in the proof of Lemma 4.1.2, this can be patched to a section σ of $\mathcal{L}_{-k}^S$, accommodating up to $m'_j a_j$ poles. Because $k \le \sum_{j=1}^e m'_j a_j + h$ we can match all the needed poles of σ, though σ might also has some simple zeros. Therefore the standard patching process gives us an immersed curve C' in class

$$B' := S + \sum m'_j E_j + \delta(B), \tag{4.1.4}$$

where $\delta(B)$ is the sum of the classes of those components of C^B that are attached to a pole of σ. Therefore $k \le \sum_{j=1}^e m'_j a_j + \delta(B) \cdot S$. Further $\delta(B) = B$ unless $B^2 = 0$, in which case $(\delta(B))^2 = 0$ as well. Hence we have

$$\begin{aligned} (B')^2 \;&\ge\; S \cdot S + (\sum m'_j E_j)^2 + 2S \cdot \delta(B) + 2S \cdot (\sum m'_j E_j) \\ &=\; -k - \sum (m'_j)^2 + 2S \cdot \delta(B) + 2 \sum m'_j a_j \\ &\ge\; -k + 2S \cdot \delta(B) + \sum m'_j a_j \;\ge 0, \end{aligned}$$

where the second inequality holds because $m'_j \le a_j$ and the third follows from (4.1.3). Hence we have a representative of A consisting of copies of S with total multiplicity $(\ell - 1)$ together with the B'-curve we have just constructed, as well as the other components of B and the E_j that did not go into making B'. By Lemma 3.2.1, we can adjoin to the B'-curve any of the components in classes B or E_j that intersect it, thereby getting a stable map Σ^A as in Proposition 3.2.4 involving copies of S with total multiplicity $(\ell - 1)$. This completes the inductive step. □

Proof of Theorem 1.2.1: general case. Now $\mathcal{S} = (C_1^S, \dots, C_s^S)$ consists of several chains of curves in classes $S_1, \dots, S_s$ with self-intersection numbers $-k_1, \dots, -k_s$. As in the special case already considered, for every $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ there is some stable map representing the class A. Moreover, we may suppose that this is $\mathcal{S}$-normalized, so that

$$A = \sum_{i=1}^{s} \ell_i S_i + \sum_{j=1}^{e} m_j E_j + B, \qquad E_j \cdot B = 0. \tag{4.1.5}$$

Our aim is to "detach" the A-stable map from $\mathcal{S}$ as much as possible to obtain an A-curve that intersects $\mathcal{S}$ in a finite number of points using the patching processes explained in Lemma 4.1.1 and 4.1.2. These are local processes that only alter the A-stable map and J near the components of $\mathcal{S}$ where patching occurs. Hence, we can forget the components of $\mathcal{S}$ with $\ell_i = 0$. We then argue by induction on the total multiplicity $\ell = \sum_{i=1}^s \ell_i$.

The process is as follows: starting with $\Sigma_A = (\Sigma^A)_0$ we construct a new stable map of the form $(\Sigma^A)_1'$ with decomposition $A = \sum_{i=1}^s \ell_i^1 S_i + \sum_j (n_j')^1 (B_j')^1$ as in (2.3.1) where $\sum_{i=1}^s \ell_i^1 < \ell := \sum_{i=1}^s \ell_i$. We then carry out the process in Proposition 3.2.4 to obtain the $\mathcal{S}$-normalized stable map $(\Sigma^A)_1$ with these multiplicities (ℓ_i^1). The new values (ℓ_i^1) are obtained from the (ℓ_i) as follows.

- We first choose any maximal connected chain $\mathcal{S}_m$ in $\mathcal{S}$.
- Next, pick a local maximum ℓ_m for the multiplicities $\ell_{i_0}, \ell_{i_0+1}, \dots, \ell_{i_0+k}$ of the elements in $\mathcal{S}_m$ and denote by $I = [i_1, \dots, i_2] \ni \{m\}$ the largest interval containing m on which ℓ_i equals this maximal value ℓ_m.
- Then there is a nonempty set I_0 that is either equal to this maximal chain I or equal to the singleton $\{i\}$ for some $i \in \mathcal{S}_m$, and we set $\ell_i^1 := \ell_i$ for $i \notin I_0$ and $\ell_i^1 := \ell_i - 1$ for $i \in I_0$.
- Finally we discard the components of $\mathcal{S}$ with new multiplicities $\ell_i^1 = 0$, obtaining a new singular set $\mathcal{S}'$.

Thus the first step is to choose a maximal connected chain in $\mathcal{S}$. If there is one consisting of a single sphere we reduce ℓ by carrying out the process described in the special case. Otherwise, choose such a chain $\mathcal{S}_m$, and label its spheres $S_i, i = 1, \dots, s_m$. For convenience, we also define $\ell_0 := 0 =: \ell_{s_m+1}$, ignoring the other spheres in $\mathcal{S}$.

For each $i = 1, \dots, s_m$ denote

$$h_i := B \cdot S_i, \quad E_j \cdot S_i = a_{ij}.$$

Because $A \cdot S_i \geq 0$ for all i and $A \cdot E_j \geq 0$ for all j we have

$$\ell_{i-1} - \ell_i k_i + \ell_{i+1} + \sum_j m_j a_{ij} + h_i \geq 0, \quad \forall i, \tag{4.1.6}$$

$$\sum_i \ell_i a_{ij} - m_j \geq 0 \quad \forall j. \tag{4.1.7}$$

The new difficulty is that we may not be able to treat each sphere in $\mathcal{S}_m$ in isolation. For example, suppose that $\mathcal{S}_m$ has two intersecting spheres each with self-intersection

-3, and that A has $\ell_1 = \ell_2 = 1$, $h_1 = h_2 = 2$ and no components E_j. Then, we can patch the B curve to $C_1^S \cup C_2^S$ in one step via Lemma 3.1.3 provided that we choose the meromorphic section σ_1 of the normal bundle to C_1^S to have poles at its intersection points with $C_2^S \cup C^B$, and similarly choose the poles of σ_2 at $C_2^S \cap (C_1^S \cup C^B)$. This gives a curve in class $S_1 + S_2 + B$ disjoint from $\mathcal{S}$. Similarly, if $\ell_1 = \ell_2 = 2$, (4.1.6) implies that $h_i \geq 4$ for $i = 1, 2$. Patching one copy of $C_1^S \cup C_2^S$ to C^B as above gives an immersed curve in class $B' = B + S_1 + S_2$ that intersects $\mathcal{S}_m$ orthogonally, because, as usual, the sections σ_i never vanish. Therefore we can repeat this process to get an immersed curve in class A that is orthogonal to $\mathcal{S}_m$. Hence its embedded resolution is the curve we are looking for. Thus, as in the case $s = 1$ considered above, the problem is essentially combinatorial. To explain the idea we first restrict to the simplest case.

Case 1: *No E_j intersects $\mathcal{S}_m$.*

In this case there are $\sum_{i=1}^{s_m} h_i$ transverse intersections of C_B with $\mathcal{S}_m$, and because we are assuming that $\mathcal{S}_m$ is connected, each interior S_i has two intersection points with $\mathcal{S}_m$, while the end curves S_1, S_{s_m} have one each. If $h_i \geq k_i$ for some i, then we can reduce ℓ_i by one by patching a section σ_i of $\mathcal{L}_{k_i}^{S_i}$ with poles at k_i distinct points in $B \cap S_i$ to B. Thus in this case we take $I_0 = \{i\}$.

From now on, we assume that $h_i < k_i$ for all i. Choose the interval $I = \{i_1 \leq i \leq i_2\}$ on which the function $i \mapsto \ell_i$ is constant and maximal. In particular $\ell_{i_1-1} < \ell_{i_1}$ and $\ell_{i_2+1} < \ell_{i_2}$. If $i_1 = i_2 = i$, then $\ell_i > \ell_{i\pm1}$. Because at least one of $\ell_{i\pm1}$ is nonzero (since $s_m > 1$), we must have $\ell_i \geq 2$. Then (4.1.6) implies that

$$2 \;\leq\; k_i \;\leq\; \frac{\ell_{i-1} + \ell_{i+1} + h_i}{\ell_i} \leq \frac{2\ell_i - 2 + h_i}{\ell_i}, \tag{4.1.8}$$

we must have $h_i \geq 2 + \ell_i(k_i - 2)$. Since $k_i \geq 2$, this implies $h_i \geq k_i$, contrary to hypothesis. Hence we must have $i_2 - i_1 \geq 1$. Then (4.1.6) implies that $k_i \leq 2 + h_i$ for $i_1 < i < i_2$. Further,

$$k_{i_1} \leq \ell_{i_1-1}/\ell_{i_1} + 1 + h_{i_1}/\ell_{i_1},$$

implies that $k_{i_1} \leq 1 + h_{i_1}$. Similarly, $k_{i_2} \leq 1 + h_{i_2}$. Therefore we can attach a curve in the class $\sum_{i=i_1}^{i_2} S_i$ to B by considering the representative of $\sum_{i=i_1}^{i_2} S_i$ formed by the union of meromorphic sections $\sigma_i, i_1 \leq i \leq i_2$, with poles either at intersection points with C^B or with the adjacent C_i^S for $i_1 \leq i \leq i_2$. In order for the resulting curve C' to be transverse to $\mathcal{S}$ it is very important that the above sections σ_i are chosen to have poles at all the intersection points of S_i with the adjacent spheres $S_{i\pm1}$ in the chain $S_{i_1}, \ldots, S_{i_2}$. For then part (ii) of Lemma 3.1.3 implies that adjacent spheres are patched pole to pole and hence by nonvanishing sections. We can choose sections with this property because, by assumption, $k_i \geq 2$ for all i. This gives a curve C' that is transverse to $\mathcal{S}$ and J-holomorphic for some $J \in \mathcal{J}(\mathcal{S})$, and hence a representative for the class A with reduced ℓ. Hence the result holds by induction.

Case 2: *The general case.*

We use the ideas in the proof of the special case $s = 1$. For each $S_i \in \mathcal{S}_m$, define

$$J(i) := \{j : E_j \cdot S_i \neq 0\},$$

and note that $a_{ij} = E_j \cdot S_i \ge 1$ for all $j \in J(i)$. The analog of condition (4.1.2) is

$$k_i \;\le\; \sum_{j\in J(i)} m'_j a_{ij} + h_i, \quad 1 \le m'_j \le \min(m_j, a_{ij}). \tag{4.1.9}$$

If there is a solution $\{m'_j : j \in J(i)\}$ to these inequalities for some i, then, using the construction in Lemma 4.1.2 for (4.1.4), we can form a curve in class $\sum_{j\in J(i)} m'_j E_j + \delta(B) + S_i$ that is orthogonal to $\mathcal{S}$, where $\delta(B)$ is the sum of the components of B that are part of this curve. Taking this together with the other components of Σ^A gives a nodal curve $(\Sigma^A)'_1$ of the form considered in Proposition 3.2.4 with $\sum_i \ell_i$ reduced by 1.

We now claim that, just as in Case 1, if there is i such that $\ell_i > \ell_{i\pm1}$, then (4.1.9) has a solution. In fact, if we replace h_i in that argument by $h'_i := \sum_{j\in J(i)} m_j a_{ij} + h_i$, then we find that $k_i \le h'_i = \sum_{j\in J(i)} m_j a_{ij} + h_i$. The proof that (4.1.2) has a solution then implies that (4.1.9) has a solution for this i. Therefore, in this case we take $I_0 = \{i\}$.

If the ℓ_i have no local strict maximum, then there is a subchain $\mathcal{S}_m := \{S_i, i_1 \le i \le i_2\}$ with $i_2 > i_1$, and $\ell_i = \ell_{i_1} > \max(\ell_{i_1-1}, \ell_{i_2+1})$ for all $i_1 \le i \le i_2$. Define:

$$\varepsilon(i) := 2, \;\; i_1 < i < i_2, \qquad \varepsilon(i_1) = \varepsilon(i_2) := 1.$$

We first claim that it is possible to find integers m'_{ij} such that for all $i_1 \le i \le i_2$ and all relevant j we have:

$$k_i \;\le\; \varepsilon(i) + \sum_{j\in J(i)} m'_{ij} a_{ij} + h_i, \quad 0 \le m'_{ij} \le \min(m_j, a_{ij}). \tag{4.1.10}$$

As in the case of the equation (4.1.2), because $a_{ij} \ge 1$ for all $j \in J(i)$, this equation has a solution provided that

$$k_i \;\le\; \varepsilon(i) + \sum_{j\in J(i)} \min(m_j, a_{ij}) a_{ij} + h_i. \tag{4.1.11}$$

Next, to solve (4.1.11), first consider the case $i = i_1$. If $\ell_{i_1} = 1$, then $i_1 = 1$ (and $i_2 = s$), and $m_j \le a_{ij}$ for all i, so that $\ell_{i_1-1} = 0$ and (4.1.11) is the same as (4.1.6). Otherwise, $\ell_{i_1} \ge 2$ and we argue as in (4.1.3): namely, taking $i = i_1$ and summing over $j \in J(i_1)$ we have

$$\begin{aligned} k_{i_1} \;&\le\; 1 + \frac{\ell_{i_1-1}}{\ell_{i_1}} + \frac{1}{\ell_{i_1}}\Big(\sum_{m_j\le a_{i_1j}} m_j a_{i_1j} + \sum_{m_j > a_{i_1j}} m_j a_{i_1j}\Big) + \frac{h_{i_1}}{\ell_{i_1}} \\ &\le\; 1 + \frac{\ell_{i_1-1}}{\ell_{i_1}} + \sum_{j\in J(i_1)} \min(m_j, a_{i_1j}) a_{i_1j} + h_{i_1}. \end{aligned}$$

Therefore (4.1.11) holds because k_{i_1} is an integer while the only term on the right that is not an integer is $\frac{\ell_{i_1-1}}{\ell_{i_1}} < 1$. A similar argument works at $i = i_2$. The argument is easier for $i_1 < i < i_2$ because then $\varepsilon(i) = 2$ and all equations, both (4.1.11) and (4.1.6) divided by ℓ_i, have an added 2 on both sides.

Now let $m''_j = \max_i m'_{ij}$. We will build a curve C' in class

$$\sum_{i_1 \le i \le i_2} S_i + \sum_j m''_j E_j + \delta(B) \tag{4.1.12}$$

in the usual way from sections σ_i over C^S_i with simple poles, multisections $(f_j)_*(\rho_j)$ over C^E_j, and some components of the curve C^B lying in the class $\delta(B)$. Further:

(a) For each j there is $i := i(j)$ such that $m''_j = m'_{ij} \le a_{ij}$. Hence we may choose f_j so that the images of the branch points of f_j form a subset of the a_{ij} disjoint points in $C^S_i \cap C^E_j$. Then we require that ρ_j has simple poles at the branch points of f_j as before.

(b) We assume that for $i \in \mathcal{S}_m$ each σ_i has poles at its $\varepsilon(i)$ intersections with the other spheres in $\mathcal{S}_m$ and other poles as described in (c), but no poles at its intersections (if any) with the spheres in $\mathcal{S} \setminus \mathcal{S}_m$.

(c) For each i and for each j with $i(j) = i$, we choose σ_i to have poles of order m''_j at each of the branch points of f_j. For j with $i \ne i(j)$, the multisection $(f_j)_*(\rho_j)$ meets C^S_i at m''_j disjoint points that are close to the intersections $C^S_i \cap C^E_j$, and we allow σ_i to have simple poles at as may of these points as necessary. Since the curves $C^E_j, j = 1, \dots, e$, are disjoint, these conditions are compatible. Moreover, the inequality (4.1.10) shows that a suitable section σ_i does exist, it might have to have to vanish at some points.

With these sections, we now construct the curve C' in class $B' = \sum_{i_1 \le i \le i_2} S_i + \sum_j m''_j E_j + \delta(B)$ by patching the $S_i \in \mathcal{S}_m$ with the multisections $(f_j)_*(\rho_j)$ and the appropriate components of C^B. Finally, we resolve all double points of C' to get an embedded curve $(C^B)'$. By construction this will be orthogonal to $\mathcal{S}$ except possibly at the zeros of the sections σ_i and the points where the graph of σ_{i_1} meets $S_{i_1 - 1}$ and the graph of σ_{i_m} meets S_{i_2+1}. As in Lemma 3.2.1, we can perturb $(C^B)'$ so that these intersections are also orthogonal. This constructs the required stable curve $(\Sigma_A)'_1$ with $\ell_1 = \ell - (i_2 - i_1 + 1) < \ell$, and hence this completes the inductive step. $\square$

Corollary 4.1.3. *Given multiplicities $\ell_i, i = 1, \dots, s$, define $A' := A - \sum_{i=1}^s \ell_i S_i$ and suppose that*

(i) $\mathrm{Gr}(A') \ne 0$,

(ii) *each $E \in \mathcal{E}$ such that $E \cdot A' < 0$ satisfies $E \cdot S_i \ge 0$ for all $i = 1, \dots, s$.*

Then A has a representing $\mathcal{S}$-normalized stable curve Σ^A with these multiplicities ℓ_i.

Proof. Consider the class A' and write it as $\sum m_j E_j + B$ where B is reduced and $B \cdot E_j = 0, E_i \cdot E_j = 0$ for $i \ne j$. Since $E_j \cdot S_i \ge 0$ for all i, j by (ii), the proof of Proposition 1.2.5 shows that each class E_j is represented by an embedded sphere for all $J \in \mathcal{J}_{SW}(\mathcal{S}, \omega, A)$. Further, (i) implies that $d(B) \ge 0$ so that we may apply Theorem 1.2.1 to conclude that for some $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ the class B is represented by a J-holomorphic embedded curve that is orthogonal to $\mathcal{S}$. Since $\mathcal{J}_{reg}(\mathcal{S}, \omega, A) \subset \mathcal{J}_{SW}(\mathcal{S}, \omega, A)$, it follows that there is $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ such that all the classes E_j, B are represented by embedded curves. Since $E_j \cdot B = 0$, these J-holomorphic curves are necessarily disjoint, and by Lemma 3.2.1 we can maneuver them to be orthogonal to $\mathcal{S}$. Therefore we may form the required nodal curve using these embedded curves. $\square$

Corollary 4.1.4. *Corollary 1.2.3 holds.*

Proof. Since the condition $g(A) = g$ is equivalent to $c_1(A) = 2 - 2g + A^2$ it is homological in nature. Hence the embedded curve C^A whose existence is guaranteed by Theorem 1.2.1 must have genus g. It is constructed from the initial nodal curve $(\Sigma^A)'$ in stages by an iterative process, starting with $(\Sigma^A)_0$ which is the cleaned up version of $(\Sigma^A)'$ constructed in Proposition 3.2.4, and then forming $(\Sigma^A)_1'$ and $(\Sigma^A)_1$, then $(\Sigma^A)_2'$ and $(\Sigma^A)_2$ and so on, as in the proof of Theorem 1.2.1. Note that the initial cleaning up process in Proposition 3.2.4 is possible only because the fact that $(B')^2 \geq 0$ allows us to arrange that B' is simple. This is why we need the initial J to be generic.

Now notice that at each step, the stable map $(\Sigma^A)_k'$ contains some of the components of $(\Sigma^A)_{k-1}$ together with a new immersed curve C' in a class B' of the form (4.1.12), which is obtained by patching (approximations to) the remaining components via Lemma 3.1.3. These "approximations" are graphs of meromorphic sections and to form C' we only resolve intersections that involve at least one pole. Then $(\Sigma^A)_k$ is constructed from $(\Sigma^A)_k'$ by first resolving intersections of C' either with itself or with the other components of $(\Sigma^A)_k' \setminus \mathcal{S}$ and then making the intersections with $\mathcal{S}$ orthogonal. At each stage the maximal genus of a component of $(\Sigma^A)_k'$ or $(\Sigma^A)_k$ can only increase, as can the sum of the genera of these components. Hence if C^A has genus g, the sum of the genera of the components of all the intermediate nodal curves $(\Sigma^A)_k'$ or $(\Sigma^A)_k$ must be $\leq g$. In particular, this must hold for the starting stable map. This proves the first claim in the corollary. The second claim follows immediately from the fact that resolving any self-intersection of a connected curve increases the genus of the curve that parametrizes it. $\square$



FIGURE 4.1.1. This example is discussed in Remark 4.1.5. The components of $\mathcal{S}$ have $S_0^2 = S_1^2 = S_2^2 = -2$ but $S_3^2 = -3$, and all intersections are shown. We take $A = 2S_0 + S_1 + S_2 + S_3 + B$. This example can be realized in the nine point blow up of $\mathbb{CP}^2$ with $S_0 = L - \sum_{i=1,3,9} E_i, S_1 = E_1 - E_2, S_2 = E_3 - E_4, S_3 = L - \sum_{i=5,6,7,8} E_i$ and $B = 2L - E_5 - E_9$. Note that$\mathcal{S}$ is negative.

Remark 4.1.5. We now explain the reason for the restrictions that we put on the singular set $\mathcal{S}$ in Theorem 1.2.1. First, we required that $k_i \geq 2$. This condition is used in the argument in Case 1 of the proof of Theorem 1.2.1 that involves equation (4.1.8). A similar difficulty with this calculation occurs whenever $\mathcal{S}$ has a component S_i (such as S_0 in Figure 4.1.1) with self-intersection $-k_i$, where k_i is less than the number of components of $\mathcal{S}$ that intersect S_i. For example, if $A = 2S_0 + S_1 + S_2 + S_3 + B$ and the pattern of intersections is as in this figure, then one can check that $A \cdot S_i \ge 0$ for all i. Hence if $d(B)$ is sufficiently large we can assume that $d(A) \ge 0$. The method above would tell us first to incorporate one C_0^S component in Σ^A. But this is impossible without involving some other C_i^S since $B \cdot S_0 = 1$ and there is no meromorphic function on C_0^S with just one simple pole. In fact we cannot glue any of these components together without gluing in a two-valued section over C_0^S. On the other hand, if we did allow this we could resolve all the singularities of A in one step. More generally, to deal with more general singular sets $\mathcal{S}$ one would have to consider branched covers of the components of $\mathcal{S}$ as well as of the exceptional divisors. This procedure should allow one to remove many, if not all, of the restrictions on $\mathcal{S}$ in Theorem 1.2.1.

4.2. **Numerical arguments.**

This section proves Proposition 1.2.7. Throughout we assume that $b_2^+ = 1$.

Given any J-holomorphic stable map $(\Sigma^A)'$ where $J \in \mathcal{J}(\mathcal{S}, A)$ consider the corresponding decomposition $A = \sum_i \ell_i S_i + \sum_{j=1}^{e'} n_j B_j'$ of (2.3.1). Order the classes B_j' (assumed distinct) so that $B_j' \in \mathcal{E}$ for $j \le p$ and $B_j' \notin \mathcal{E}$ otherwise, and write $E_j' := B_j'$ for $1 \le j \le p$, and $B' := \sum_{j>p} n_j' B_j'$. Then

$$A \;=\; \sum_i \ell_i S_i + \sum_{j=1}^{p} n_j' E_j' + \sum_{j>p} n_j' B_j' \;=\; \sum_i \ell_i S_i + \sum_{j=1}^{p} n_j' E_j' + B'. \tag{4.2.1}$$

We will assume that the decomposition is **nontrivial**, in the sense that $A \ne B'$. When A is reduced this is equivalent to requiring that some $\ell_i > 0$. Note that the above decomposition is not the same as that in (3.2.2) since we may have $B' \cdot E_j' \ne 0$ for some $j \le p$. Further,

(i) $B' \ne 0$ implies $B' \cdot (A - B') > 0$

because $(\Sigma^A)'$ is connected. Moreover, if $J \in \mathcal{J}_{reg}(\mathcal{S}, A)$, then Lemma 2.2.4 (ii) implies that we also have

(ii) $(B_j')^2 \ge 0$ for $j > p$.

Lemma 4.2.1. *Suppose that $b_2^+ = 1$, let $\mathcal{S}$ be some singular set whose components are spheres, and suppose that A satisfies the conditions of Theorem 1.2.1. Suppose further that for any decomposition* (4.2.1) *for A that satisfies (ii) above we have $d(B') < d(A)$. Let $k_A := \frac{1}{2} d(A)$. Then, for $J \in \mathcal{J}_{reg}(\mathcal{S})$, there is an embedded J-holomorphic A-curve through every generic k_A-tuple of points in M.*

Proof. By definition of $\mathrm{Gr}(A)$ there is for each ω-tame J a J-holomorphic representative of the class A through every set of $k_A := \frac{1}{2} d(A)$ points in M. Let $J \in \mathcal{J}_{reg}(\mathcal{S}, \omega, A)$ and consider a J-holomorphic representative $(\Sigma^A)'$ of A with nontrivial decomposition (4.2.1). If we remove the rigid components in the classes $\ell_i S_i$ and $n_j' E_j'$ from $(\Sigma^A)'$ we are left with a (possibly disconnected) stable map Σ' in the class $B' := \sum_{j>p} n_j' B_j'$, where by construction $B_j', j > p$, are the classes of its underlying simple components. Since J is generic, and $d(B_j') := (B_j')^2 + c_1(B_j')$ is the Fredholm degree for simple curves in class B_j', each such component has $d(B_j') \ge 0$, and hence $(B_j')^2 \ge 0$ by Lemma 2.2.4 (ii). Therefore condition (ii) holds, so that $d(B') < d(A)$ by hypothesis.

Suppose that $(B_j')^2 = 0$ for $j \in K_1$ and $(B_j')^2 > 0$ for $j \in K_2$. If $(B_j')^2 > 0$ then the corresponding component in $(\Sigma^A)'$ is a multiply covered copy of a simple B_j'-curve and cannot go through more than $\frac{1}{2}d(B_j')$ generic points. Moreover $d(n_j'B_j') \ge d(B_j')$. On the other hand, if $(B_j')^2 = 0$ then the corresponding components in $(\Sigma^A)'$ go through at most $d(n_j'B_j') = n_j' d(B_j')$ generic points. Hence the nodal curve $(\Sigma^A)'$ can go through at most $\frac{1}{2}D$ generic points, where

$$D := \sum_{j \in K_1} n_j' d(B_j') + \sum_{j \in K_2} d(B_j').$$

But because $B_j' \cdot B_k' \ge 0$ for all $j \ne k$ it is easy to check that $D \le d(B')$. But $d(B') < d(A) = 2k_A$. Therefore, by standard arguments, we can choose k_A points that do not lie on any such nodal curve $(\Sigma_A)'$. □

We now give two different sets of conditions under which $d(B') < d(A)$. First we discuss the genus 0 situation, using an argument adapted from Li–Zhang [LZ12, Lemma 4.9].[5]

Lemma 4.2.2. *Suppose that $b_2^+ = 1$, let $\mathcal{S}$ be any singular set, and suppose that A, satisfying the conditions of Theorem 1.2.1, has $g(A) := 1 + \frac{1}{2}(A^2 - c_1(A)) = 0$ and $d(A) := A^2 + c_1(A) > 0$. Then for any nontrivial decomposition* (4.2.1) *for A that satisfies condition (ii) we have $d(B') < d(A)$.*

Proof. The statement holds when $B' = 0$ since $d(0) = 0 < d(A)$. Therefore we may suppose that $B' \ne 0, A$. Next notice that for any class B with a connected simple J-holomorphic representative, we have $g(B) = 1 + \frac{1}{2}(B^2 - c_1(B))$ so that $\frac{1}{2}d(B) = 1 + B^2 - g(B)$. If there are any B_j' in (4.2.1) with $(B_j')^2 > 0$ or if there are two classes B_j' of zero square, then by Lemma 2.1.1 the B' curve is connected, and hence has a simple representative for some J by Lemma 3.2.1. Therefore, $\frac{1}{2}d(A) = 1 + A^2$ while $\frac{1}{2}d(B') \le 1 + (B')^2$. Therefore, it suffices to show that $(B')^2 < A^2$. But

$$A^2 - (B')^2 = (A+B')\cdot(A-B') = A\cdot\Big(\sum \ell_i S_i + \sum n_j' E_j'\Big) + B'\cdot(A-B') \ge B'\cdot(A-B') > 0,$$

where the first inequality holds because of the assumptions on A in Theorem 1.2.1 and the second (strict) inequality holds because, as we noted above, $(\Sigma^A)'$ is connected and $A \ne B'$.

This completes the proof unless $B' = nB_0'$ where $(B_0')^2 = 0$. In this case, by Lemma 2.2.4 (ii) either $g(B_0') = 1$ and $d(B_0') = d(nB_0') = 0$, or $g(B_0') = 0$ and

---

[5] Instead of condition (ii) they use the hypothesis that A is J-NEF, which implies that $A \cdot B_j' \ge 0$ for all j.

$d(nB_0') = n$. In the first case we still have $\frac{1}{2}d(B') \le 1 + (B')^2$, and so the previous argument applies. It also applies if $n = 1$ and $g(B_0') = 0$. Therefore, it remains to consider the case $g(B_0') = 0$ and $d(nB_0') = n > 1$. Then $B' \cdot (A - B') \ge n$ since B' has n disjoint components, while $\frac{1}{2}d(B') = \frac{n}{2} \le n - 1$. Hence $d(B') < d(A)$ if $n - 1 < A^2 = A^2 - (B')^2$. But this holds by the calculation above and the fact that $B' \cdot (A - B') > n - 1$. □

Secondly we extend an argument from Biran [B99]. Note that it does not require $b_2^+ = 1$ or condition (ii). Instead, we have a strong condition on $\mathcal{S}$.

Lemma 4.2.3. *Let $\mathcal{S}$ be any singular set whose components satisfy $c_1(S_i) = 0$ for all i, and let A satisfy the conditions of Theorem 1.2.1. Then for any decomposition* (4.2.1) *for A, that is nontrivial in the sense that $\sum_j E_j' + B' \ne 0$, we have $d(B') < d(A)$.*

Proof. Let us write $Z := \sum_i \ell_i S_i$. First suppose that $B' = 0$ so that $A = Z + \sum_j n_j' E_j'$. We must show that $d(A) > 0 = d(B')$. By assumption $A \ne Z$. Further,

$$d(A) = A^2 + c_1(Z + \sum_j n_j' E_j') = A^2 + \sum n_j' > 0$$

unless $A \in \mathcal{E}$ and $\sum n_j' = 1$. But then $A = Z + E$ where $E \in \mathcal{E}$, so that

$$0 = A^2 + 1 = Z^2 + 2Z \cdot E + E^2 + 1 = Z^2 + 2Z \cdot E.$$

On the other hand, by assumption $A \cdot Z = Z^2 + E \cdot Z \ge 0$. Since $Z \cdot E > 0$, this is impossible. Hence when $B' = 0$ we have $d(A) > 0$ as required.

Now suppose that $B' \ne 0$. Note also that $A \ne E_j$ for any j with $m_j > 0$ since $\omega(A - E_j) > 0$. Hence

$$A \cdot E_j \;=\; \big(Z + \sum_k m_k E_k\big) \cdot E_j \;=\; Z \cdot E_j - m_j \;\ge\; 0$$

so that

$$0 \;\le\; m_j \;\le\; Z \cdot E_j, \quad \forall j. \tag{4.2.2}$$

By assumption on the S_i, we have $c_1(Z) = 0$. Therefore

$$\begin{aligned} d(A) - d(B') \;&=\; \big(Z + \sum_j n_j' E_j' + B'\big) \cdot A + \sum_j n_j' c_1(E_j') + c_1(B') - d(B') \\ &=\; \big(Z + \sum_j n_j' E_j' + B'\big) \cdot A + \sum_j n_j' c_1(E_j') - (B')^2 \\ &=\; Z \cdot A + \sum_j n_j' E_j' \cdot A + B' \cdot (A - B') + \sum_j n_j' c_1(E_j') \\ &> Z \cdot A + \sum_j n_j' \ge 0, \end{aligned}$$

where the strict inequality uses the fact that $E_j' \cdot A \ge 0$ and that $B' \cdot (A - B') > 0$. This completes the proof. □

Corollary 4.2.4. *Proposition 1.2.7 holds.*

Proof. Part (i) is an immediate consequence of Lemmas 4.2.3 and 4.2.1. Part (ii) holds by Lemma 1.1.3. Finally, the last claim holds when $d(A) = 0$ by Proposition 1.2.5 (because $d(A) = g(A) = 0$ for a reduced class implies that $A \in \mathcal{E}$), and holds when $d(A) > 0$ by Lemmas 4.2.2 and 4.2.1. $\square$

Remark 4.2.5. (i) Biran actually assumed that $A \cdot S_i + c_1(S_i) \geq 0$ for all i. However, In the applications of interest to us, we usually know only that $A \cdot S_i \geq 0$. Hence we replaced Biran's condition by $c_1(S_i) = 0$ for all i.

(ii) We assumed in Lemma 4.2.1 that the components of $\mathcal{S}$ are spheres for the following reason. The crucial condition that makes the calculation in Lemma 4.2.3 work is that $c_1(S_i) = 0$ for all i. If $g(S_i) > 0$ then because $c_1(S_i) = 2 - 2g(S_i) + S_i^2$ we would either have $g(S_i) = 1$ and $S_i^2 = 0$ or $S_i^2 > 0$. In the second case $d(S_i) = c_1(S_i) + S_i^2 > 0$, so that the S_i-curve would not be rigid, and the argument in Corollary 4.2.4 would fail. In the first case, the argument would hold but would tell us nothing new because for generic elements $J \in \mathcal{J}(\mathcal{S})$ the component S_i is regular. Hence we could reduce to the case when all components of $\mathcal{S}$ have genus 0 by simply removing these regular components S_i.

(iii) The statement of Proposition 1.2.7 refers only to the existence of a single embedded representative of A for a single generic J. However, the above proofs can easily be modified to get existence for generic 1-parameter families J_t (as needed for inflation, cf. Lemma 4.3.3): the arguments in Lemmas 2.2.4 (ii) and 4.2.1 both hold for 1-parameter families of J since the index $\mathrm{Ind} D_{u,J_t}$ is always even. (Cf. the proof of Lemma 2.2.3 (iii).) This modification of part (i) of Proposition 1.2.7 is therefore all that is needed to complete the proof of the key technical Lemma 2.18 in [Mc09ii], since in that paper all the curves in $\mathcal{S}$ have self-intersection -2.

4.3. **Symplectic inflation.** We now prove Proposition 1.2.9. Throughout this section we assume that (M, ω) is a blow up of a rational or ruled manifold. The proof is an adaptation of the inflation method that was developed by Lalonde[L94] and [Mc98], and then extended by Biran [B99, §6]. When $\mathcal{S}$ is empty, the key step is to prove the following claim.

Lemma 4.3.1 ([Mc98], Lemma 1.1)**.** *Let A be a reduced class in $H_2(M;\mathbb{Z})$ with non-negative self-intersection number and nonzero Gromov invariant* $\mathrm{Gr}(A)$. *Then, given any smooth family $\omega_t, 0 \leq t \leq 1$, of symplectic forms on M with $\omega_0 = \omega$, there is (after possibly reparametrizing with respect to t) a family of closed forms ρ_t on M in the class Poincaré dual to A such that the family*

$$\tau_{\kappa t} := \omega_t + \kappa\rho_t, \quad 0 \leq t \leq 1,$$

is symplectic for all $\kappa \geq 0$.

The statement in Proposition 1.2.9 (for $\mathcal{S} = \emptyset$) then follows by carefully choosing classes A along which to perturb $[\omega_t]$. By this means we construct a 2-parameter family $\omega_{st} := \omega_t + \kappa_s\rho_t, 0 \leq s, t \leq 1$, of symplectic forms such that

- $\omega_{0t} = \omega_t$ for $t \in [0,1]$;
- $\omega_{s0} = \omega$ and $\omega_{s1} = \omega'$ for all s;
- $[\omega_{1t}] = [\omega]$ for all t.

Then the isotopy ϕ_t exists by applying Moser's argument to the family $\omega_{1t}, 0 \le t \le 1$.

When $\mathcal{S}$ is nonempty, we use the same strategy of proof. In fact the arguments in [Mc98] prove the following.

Lemma 4.3.2. *Suppose that for each family ω_t satisfying the conditions of Proposition 1.2.9 and for each reduced class A in $H_2(M;\mathbb{Z})$ with $A \in \mathcal{P}^+$ and $A \cdot S_i \ge 0$ for all i, there is (after reparametrization) a family of closed forms $\rho_t, t \in [0,1]$, on M such that for all $t \in [0,1]$*

(i) *$[\rho_t]$ is Poincaré dual to A;*
(ii) *$\rho_t|_{S_i} \ge 0$ for all i and has support disjoint from the intersections $C_i^S \cdot C_j^S$; and*
(iii) *the forms $\tau_{\kappa t} := \omega_t + \kappa\rho_t$ are symplectic for all $\kappa \ge 0$.*

Then Proposition 1.2.9 holds.

The first step in the proof of Lemma 4.3.1 is to notice that when $A \in \mathcal{P}^+$ we have $d(qA) \to \infty$ as $q \to \infty$ because q^2A^2 grows quadratically with q while $K \cdot qA$ grows linearly. Hence Lemma 2.1.3 implies that $\mathrm{Gr}(qA) \ne 0$ for some $q > 0$. One then shows that for any generic smooth family of ω_t-tame almost complex structures J_t, there is a corresponding family Z_t of embedded J_t-holomorphic curves in class qA. This may not hold for an arbitrary generic path in $\mathcal{J}(\mathcal{S})$. Instead, we prove the following slightly more complicated statement.

Lemma 4.3.3. *Let $\omega_t, t \in [0,1]$, be a path of symplectic forms as in Proposition 1.2.9, and suppose given a reduced class $A \in \mathcal{P}^+$ such that $A \cdot S_i \ge 0$ for all i. Suppose further that $d(A) \ge 0$ (in the rational case) or that $d(A) \ge g + \frac{k}{4}$ (if M is the k-point blow up of a ruled surface). Choose a smooth path $J_t \in \mathcal{J}(\mathcal{S}, \omega_t, A), t \in [0,1]$. Then, possibly after reparametrization with respect to t, $(J_t)_{t\in[0,1]}$ can be perturbed to a smooth path $\big(J'_t \in \mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)\big)_{t\in[0,1]}$ such that there is a smooth family $C_t^A, t \in [0,1]$, of embedded J'_t-holomorphic curves in class A.*

Proof. Since the space $\bigcup_{t\in[0,1]} \mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)$ is path connected by Lemma 2.2.3, we may choose (J_t) to be a generic path in $J_t \in \mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)$ for all t. Consider the following compact space of stable maps:

$$X := \bigcup_{t\in[0,1]} \overline{\mathcal{M}}(A, J_t).$$

This space is stratified according to the topological type T of the domains of the stable maps, where T keeps track both of the structure of the domain and the homology classes of the corresponding curves. These strata X_T are ordered by the relation that $T' \le T$ if a stable with domain of type T can degenerate into one of type T'. Since J_t ranges in a compact set there are a finite number of such decompositions $A = \sum_i \ell_i S_i + \sum_j m'_j B'_j$ as in (3.2.1). If we "clean up" each of the corresponding nodal curves $(\Sigma^A)'$ to Σ^A with decomposition $A = \sum_{i=1}^{s} \ell_i S_i + \sum_{j=1}^{e} m_j E_j + B$ as in Proposition 3.2.4, then there still are only a finite number of such cleaned up decompositions, since the classes B, E_j are obtained from the B_j' by an essentially homological process. Let $d_{\max}$ be the maximum of the numbers $d(B)$, where B occurs in such a decomposition. Note that $d_{\max} \geq d(A)$ since we could have $\ell_i = m_j = 0$ for all i, j. Next, fix a decomposition

$$A = \sum_{i=1}^{s} \ell_i S_i + \sum_{j=1}^{e} m_j E_j + B \tag{4.3.1}$$

of this type with $d(B) = d_{\max}$ and with maximal multiplicities (ℓ_i), in the sense that there is no other representative of A with decomposition $\sum_{i=1}^{s} \ell_i' S_i + \sum_{j=1}^{e'} m_j' E_j' + B'$ where $d(B') = d_{\max}$, $\ell_i' \geq \ell_i$ for all i and $\ell_i' > \ell_i$ for some i.

We next claim that $\mathrm{Gr}(B) \neq 0$. If M is rational, this is immediate from Lemma 2.1.3 (i) since $B^2 \geq 0$, $\omega(B) > 0$ by construction, and $d(B) \geq d(A) \geq 0$. So suppose that M is the blowup of a ruled surface. If $B^2 = 0$ then, by Proposition 3.2.4, we may write $B = mB_0$ where $m \geq 1$ and B_0 is represented by an embedded sphere or torus. In the first case $\mathrm{Gr}(B) = 1$ since for generic J there is a unique B-curve through each set of m generic points, while in the second $d(B_0) = d(B) = 0$. But this is impossible since $d(B) = d_{\max} \geq d(A) > 0$ by hypothesis. Therefore, it remains to consider the case when $B^2 > 0$. Since $\omega(B) > 0$ by construction, this means that $B \in \mathcal{P}^+$. Therefore $\mathrm{Gr}(B) \neq 0$ by Lemma 2.1.3 (ii) applied to the class B.

Claim: *Given A as in* (4.3.1)*, there is, after possible reparametrization with respect to t, a corresponding family $\Sigma_t^A, t \in [0,1]$, of $\mathcal{S}$-normalized J_t-holomorphic nodal curves.*

Since the classes E_j, B in (4.3.1) have nontrivial Gromov invariant, they are always represented in some form for each J_t. By Proposition 1.2.5 the classes E_j are in fact always represented by embedded curves C_{jt}^E since they lie in $\mathcal{E}$ and $J_t \in \mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)$ for all t. We next check that we can choose the path J_t so that the class B does not decompose. By standard theory (sketched below, but cf. [B99, §6]), this will follow if we can show that for each decomposition $B = \sum_j B_j'$ of the B-curve, the sum of the Fredholm indices of its nonrigid components is strictly less than the Fredholm index $d(B)$ of the class B. If the components of the B curve are all transverse to $\mathcal{S}$, then this calculation is standard. On the other hand, if for some J_t the decomposition is a stable map $(\Sigma_B)'$ that involves some components of $\mathcal{S}$, then Lemma 3.2.5 implies that this sum is at most $d(B_0)$ where B_0 is the nonrigid component of the *cleaned up* version of $(\Sigma_B)'$. But the maximality of the pair $d(B) = d_{\max}$ and (ℓ_i) implies that $d(B_0) < d(B)$, and since $d(D)$ is always even, we actually have $d(B_0) \leq d(B) - 2$. Therefore in a generic path J_t, the dimension of the moduli space of these stable maps is at most $d(B) - 1$, and hence those curves cannot go through $k := \frac{1}{2} d(B)$ generic points. Thus, the space of embedded B-curves that are J_t-holomorphic for some t and go though k generic points is a compact 1-manifold with boundary. Moreover, because $\mathrm{Gr}(B) \neq 0$ there is at least one connected component of this 1-manifold with one end at $t = 0$ and the other at $t = 1$. Taking such a component, and reparametrizing with respect to t as necessary, we therefore have a family C_t^B of embedded J_t-holomorphic curves in class B.

Next observe that the curves C_{jt}^E and C_t^B are mutually disjoint for each t and that their intersections with $\mathcal{S}$ are fixed by the homological intersection numbers $E_j \cdot S_i$ and $B \cdot S_i$. Moreover, since we are only dealing with a 1-parameter path we can assume that all intersections are transverse. (For example, if some C_{jt}^E happens to be tangent to $\mathcal{S}$, we can slightly perturb the path J_t so that this intersection is transverse since such tangencies happen in codimension at least 2.) Finally we straighten up the intersections of the E_j and B curves with $\mathcal{S}$ to make them orthogonal, using the 1-parameter version of Lemma 3.2.1 as in Remark 3.2.3. This proves the Claim.

The next step is to check that the procedure described in the proof of Theorem 1.2.1 that constructs an embedded curve C_t^A from the nodal curve Σ_t^A can be carried out smoothly as t varies in $[0,1]$. This holds because this procedure is determined by the intersection pattern of the components of the nodal curves Σ_t^A, which as we have noted is independent of t; cf. Remark 3.2.3. Since this procedure involves patching, the smooth family (J_t) changes to a smooth family (J_t'). As in Remark 3.1.4, we may assume that (J_t') is $\mathcal{C}^1$-close to (J_t). However, because $\mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)$ is not C^1-open, this by itself does not guarantee that $J_t' \in \mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)$ for all t. On the other hand, if C_t^A is a sphere (for one and hence all t), then it is automatically regular. Hence, we can perturb J_t' to make it lie in $\mathcal{J}_{SW}(\mathcal{S}, \omega_t, A)$, without changing it on $\mathcal{S}$ or on the tangent bundle TC_t^A as in Lemma 2.2.4 (i). This completes the proof. □

Corollary 4.3.4. *Given ω_t and A as in Lemma 4.3.3, we can reparametrize with respect to t so that there is a family of closed forms $\rho_t, t \in [0,1]$, on M such that for all $t \in [0,1]$*

- *$[\rho_t]$ is Poincaré dual to A;*
- *the forms $\tau_{\kappa t} := \omega_t + \kappa\rho_t$ are symplectic and adapted to $\mathcal{S}$ for all $\kappa \geq 0$.*

Proof. The required forms ρ_t can be constructed by hand as in [Mc98] to have support in an arbitrarily small neighborhood of the curves C_t^A. If $A \cdot S_i = 0$ for all i, the curve C_t^A is disjoint from $\mathcal{S}$, so that we can arrange that the forms ρ_t vanish on $\mathcal{S}$. Moreover because C^A is orthogonal to $\mathcal{S}$ it does not go through any of the intersection points $C_i^S \cdot C_j^S$ so that we can arrange that $\rho_t = 0$ near these points. so that Otherwise, one can use the method in either [B99, §6] [B99, §6] or [Mc01, Lemma 3.1] to arrange that the curves $\mathcal{S}$ are symplectically embedded for all $\kappa(t)$. □

Proof of Proposition 1.2.9. Given any A as in Lemma 4.3.2 we have already noted that qA satisfies the additional hypotheses of Lemma 4.3.3 for large $q > 0$. Hence, the hypothesis of Lemma 4.3.2 holds by Corollary 4.3.4. The methods in [Mc98] then produce an isotopy $\omega_t, 0 \leq t \leq 1$, between ω and ω'. Each form ω_t is adapted to $\mathcal{S}$ by condition (ii) of Lemma 4.3.2. Hence a standard Moser argument allows us to modify these forms near $\mathcal{S}$ so that they are constant there. Then the required isotopy clearly exists. □

References

[B99] P. Biran, A stability property of symplectic packing, *Invent. Math.* **136** (1999), 123–155.

[EKS84] A. L. Edmonds, R. S. Kulkarni and R. E. Stong, Realizability of branched coverings of surfaces, *Trans. Amer. Math. Soc.* **282** (1984) 773–790.

[H91] A. Hurwitz, Über Riemannischen Flächen mit gegebenen Verzweigungspunkten, *Math. Ann.* **103** (1981), 1–60.

[KM94] P. Kronheimer and T. Mrowka, The genus of embedded surfaces in the projective plane, *Math. Res. Letters*, (1994), 797–808

[L94] F. Lalonde, Isotopy of symplectic balls, Gromov's radius and the structure of ruled symplectic 4-manifolds, *Math. Ann.* **300** (1994), 273-296.

[LL95] Tian-Jun Li and A. Liu, General wall crossing formula, *Math Research Letters* **2** (1995), 797–810.

[LL99] Tian-Jun Li and A. Liu, On the equivalence between SW and Gr in the case $b^+ = 1$, *Internat. Math. Res. Not.* **7** (1999), 335–345.

[LL01] Tian-Jun Li and A. Liu, Uniqueness of symplectic canonical class, surface cone and symplectic cone of manifolds with $b^1 = 1$, *J. Differ. Geom.* **58** (2) (2001), 331–370.

[LZ12] Tian-Jun Li and Weiyi Zhang, J-holomorphic curves in a NEF class, arXiv:12103337

[Mc91] D. McDuff, The Local Behavior of Holomorphic curves in almost complex 4-manifolds, *Journ. Diff. Geo.*, 34 (1991), 143-164.

[Mc94i] D. McDuff, Singularities and positivity of intersections of J-holomorphic curves, pp. 191–216 in *Holomorphic curves in symplectic geometry*, ed. Audin and Lafontaine, Progress in Mathematics **117**, (1994) Birkhäuser, Basel.

[Mc94ii] D. McDuff, Notes on ruled symplectic 4-manifolds. *Transactions of the American Mathematical Society* **345** (1994) 623–639.

[Mc97] D. McDuff, Lectures on Gromov invariants, in *Gauge Theory and Symplectic Geometry* ed Hurtubise and Lalonde, NATO ASI Series, Kluwer, Dordrecht/Boston (1997).

[Mc98] D. McDuff, From Symplectic deformation to isotopy, in *Proceedings of Irvine Conference 1996*, ed Stern. International Press. (1998).

[Mc01] D. McDuff, Symplectomorphism Groups and Almost Complex Structures, *Enseignement Math.*, **38** (2001), 1–30.

[Mc09i] D. McDuff, Symplectic embeddings of 4-dimensional ellipsoids, *Journ. of Topology* (2009) **2**, 1-22

[Mc09ii] D. McDuff, Some 6 dimensional Hamiltonian S^1-manifolds, *Journ. of Topology* (2009) **2**, 589–623.

[MS96] D. McDuff and D.A. Salamon, A survey of symplectic manifolds with $b^+ = 1$, *Turkish Journal of Mathematics,* 20 (1996) 47-60.

[MS04] D. McDuff and D.A. Salamon, *J-holomorphic curves and symplectic topology.* Colloquium Publications **52**, American Mathematical Society, Providence, RI, (2004), 2nd ed. (2012).

[Tau95] C. H. Taubes, The Seiberg–Witten and the Gromov invariants, *Math. Research Letters* **2**, (1995), 221–238.

[Tau96] C. H. Taubes, Counting pseudoholomorphic submanifolds in dimension 4, *Journ Diff Geo.* **44**, (1996), 818–893.

DEPARTMENT OF MATHEMATICS, BARNARD COLLEGE, COLUMBIA UNIVERSITY NEW YORK, USA
E-mail address: `dusa@math.columbia.edu`